




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 14/00</b>		<b>A2</b>	(11) International Publication Number: <b>WO 99/33869</b>
			(43) International Publication Date: <b>8 July 1999 (08.07.99)</b>
(21) International Application Number: <b>PCT/US98/27416</b>		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: <b>22 December 1998 (22.12.98)</b>			
(30) Priority Data:			
08/998,253	24 December 1997 (24.12.97)	US	
08/998,255	24 December 1997 (24.12.97)	US	
09/118,627	17 July 1998 (17.07.98)	US	
09/118,554	17 July 1998 (17.07.98)	US	
(71) Applicant: CORIXA CORPORATION [US/US]; Suite 200, 1124 Columbia Street, Seattle, WA 98104 (US).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors: REED, Steven, G.; 2843 - 122nd Place Northeast, Bellevue, WA 98005 (US). XU, Jiangchun; 15805 South-east 43rd Place, Bellevue, WA 98006 (US).			
(74) Agents: MAKI, David, J. et al.; Seed and Berry LLP, 6300 Columbia Center, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).			

(54) Title: COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

## (57) Abstract

Compounds and methods for the treatment and diagnosis of breast cancer are provided. The inventive compounds include polypeptides containing at least a portion of a breast tumor protein. Vaccines and pharmaceutical compositions for immunotherapy of breast cancer comprising such polypeptides, or polynucleotide molecules encoding such polypeptides, are also provided, together with polynucleotide molecules for preparing the inventive polypeptides.



# **FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

### TECHNICAL FIELD

The present invention relates generally to compositions and methods for the treatment and diagnosis of breast cancer. The invention is more particularly related to polypeptides comprising at least a portion of a protein that is preferentially expressed in breast tumor tissue and to polynucleotide molecules encoding such polypeptides. Such polypeptides may be used in vaccines and pharmaceutical compositions for treatment of breast cancer. Additionally such polypeptides and polynucleotides may be used in the immunodiagnosis of breast cancer.

### BACKGROUND OF THE INVENTION

Breast cancer is a significant health problem for women in the United States and throughout the world. Although advances have been made in detection and treatment of the disease, breast cancer remains the second leading cause of cancer-related deaths in women, affecting more than 180,000 women in the United States each year. For women in North America, the life-time odds of getting breast cancer are now one in eight.

No vaccine or other universally successful method for the prevention or treatment of breast cancer is currently available. Management of the disease currently relies on a combination of early diagnosis (through routine breast screening procedures) and aggressive treatment, which may include one or more of a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone therapy. The course of treatment for a particular breast cancer is often selected based on a variety of prognostic parameters, including an analysis of specific tumor markers. See, e.g., Porter-Jordan and Lippman, *Breast Cancer* 8:73-100 (1994). However, the use of established markers often leads to a result that is difficult to interpret, and the high mortality observed in breast cancer patients indicates that improvements are needed in the treatment, diagnosis and prevention of the disease.

Accordingly, there is a need in the art for improved methods for therapy and diagnosis of breast cancer. The present invention fulfills these needs and further provides other related advantages.

## SUMMARY OF THE INVENTION

The present invention provides compounds and methods for immunotherapy of breast cancer. In one aspect, isolated polypeptides are provided comprising at least an immunogenic portion of a breast tumor protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein the breast tumor protein comprises an amino acid sequence encoded by a polynucleotide molecule having a partial sequence selected from the group consisting of (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52 and 55-67, 72, 73, and 89-94, (b) complements of said nucleotide sequences and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

In related aspects, isolated polynucleotide molecules encoding the above polypeptides are provided. In specific embodiments, such polynucleotide molecules have partial sequences provided in SEQ ID NOS: 3, 10, 17, 24, 45-52 and 55-67, 72, 73, and 89-94. The present invention further provides expression vectors comprising the above polynucleotide molecules and host cells transformed or transfected with such expression vectors. In preferred embodiments, the host cells are selected from the group consisting of *E. coli*, yeast and mammalian cells.

In another aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, an inventive polypeptide and a known breast antigen.

The present invention also provides pharmaceutical compositions comprising at least one of the above polypeptides, or a polynucleotide molecule encoding such a polypeptide, and a physiologically acceptable carrier, together with vaccines comprising at least one or more such polypeptide or polynucleotide molecule in combination with a non-specific immune response enhancer. Pharmaceutical compositions and vaccines comprising one or more of the above fusion proteins are also provided.



In related aspects, pharmaceutical compositions for the treatment of breast cancer comprising at least one polypeptides and a physiologically acceptable carrier are provided, wherein the polypeptide comprises an immunogenic portion of a breast tumor protein or a variant thereof, the breast tumor protein being encoded by a polynucleotide molecule having a partial sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88, (b) complements of said nucleotide sequences, and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions. The invention also provides vaccines for the treatment of breast cancer comprising such polypeptides in combination with a non-specific immune response enhancer, together with pharmaceutical compositions and vaccines comprising at least one polynucleotide molecule having a partial sequence provided in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88.

In yet another aspect, methods are provided for inhibiting the development of breast cancer in a patient, comprising administering an effective amount of at least one of the above pharmaceutical compositions and/or vaccines.

The present invention also provides methods for immunodiagnosis of breast cancer, together with kits for use in such methods. In one specific aspect of the present invention, methods are provided for detecting breast cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the inventive polypeptides; and (b) detecting in the sample a protein or polypeptide that binds to the binding agent. In preferred embodiments, the binding agent is an antibody, most preferably a monoclonal antibody.

In related aspects, methods are provided for monitoring the progression of breast cancer in a patient, comprising: (a) contacting a biological sample obtained from a patient with a binding agent that is capable of binding to one of the above polypeptides; (b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent; (c) repeating steps (a) and (b); and comparing the amounts of polypeptide detected in steps (b) and (c).

Within related aspects, the present invention provides antibodies, preferably monoclonal antibodies, that bind to the inventive polypeptides, as well as diagnostic kits

comprising such antibodies, and methods of using such antibodies to inhibit the development of breast cancer.

The present invention further provides methods for detecting breast cancer comprising: (a) obtaining a biological sample from a patient; (b) contacting the sample with a first and a second oligonucleotide primer in a polymerase chain reaction, at least one of the oligonucleotide primers being specific for a DNA molecule that encodes one of the above polypeptides; and (c) detecting in the sample a DNA sequence that amplifies in the presence of the first and second oligonucleotide primers. In a preferred embodiment, at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID NOS: 1-94.

In a further aspect, the present invention provides a method for detecting breast cancer in a patient comprising: (a) obtaining a biological sample from the patient; (b) contacting the sample with an oligonucleotide probe specific for a polynucleotide molecule that encodes one of the above polypeptides; and (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe. Preferably, the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a DNA molecule having a partial sequence selected from the group consisting of SEQ ID NOS: 1-94.

In related aspects, diagnostic kits comprising the above oligonucleotide probes or primers are provided.

These and other aspects of the present invention will become apparent upon reference to the following detailed description. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A and B show the specific lytic activity of a first and a second B511S-specific CTL clone, respectively, measured on autologous LCL transduced with B511s (filled squares) or HLA-A3 (open squares).

## DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the immunotherapy and diagnosis of breast cancer. The inventive compositions are generally isolated polypeptides that comprise at least a portion of a breast tumor protein. Also included within the present invention are molecules (such as an antibody or fragment thereof) that bind to the inventive polypeptides. Such molecules are referred to herein as "binding agents."

In particular, the subject invention discloses polypeptides comprising at least a portion of a human breast tumor protein, or a variant thereof, wherein the breast tumor protein includes an amino acid sequence encoded by a polynucleotide molecule including a sequence selected from the group consisting of: nucleotide sequences recited in SEQ ID NOS: 1- 94, the complements of said nucleotide sequences, and variants thereof. As used herein, the term "polypeptide" encompasses amino acid chains of any length, including full length proteins, wherein the amino acid residues are linked by covalent peptide bonds. Thus, a polypeptide comprising a portion of one of the above breast proteins may consist entirely of the portion, or the portion may be present within a larger polypeptide that contains additional sequences. The additional sequences may be derived from the native protein or may be heterologous, and such sequences may be immunoreactive and/or antigenic.

As used herein, an "immunogenic portion" of a human breast tumor protein is a portion that is capable of eliciting an immune response in a patient inflicted with breast cancer and as such binds to antibodies present within sera from a breast cancer patient. Such immunogenic portions generally comprise at least about 5 amino acid residues, more preferably at least about 10, and most preferably at least about 20 amino acid residues. Immunogenic portions of the proteins described herein may be identified in antibody binding assays. Such assays may generally be performed using any of a variety of means known to those of ordinary skill in the art, as described, for example, in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988. For example, a polypeptide may be immobilized on a solid support (as described below) and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, <sup>125</sup>I-labeled Protein A. Alternatively, a polypeptide may be used to generate

monoclonal and polyclonal antibodies for use in detection of the polypeptide in blood or other fluids of breast cancer patients. Methods for preparing and identifying immunogenic portions of antigens of known sequence are well known in the art and include those summarized in Paul, *Fundamental Immunology*, 3<sup>rd</sup> ed., Raven Press, 1993, pp. 243-247.

The term "polynucleotide(s)," as used herein, means a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases and includes DNA and corresponding RNA molecules, including HnRNA and mRNA molecules, both sense and anti-sense strands, and comprehends cDNA, genomic DNA and recombinant DNA, as well as wholly or partially synthesized polynucleotides. An HnRNA molecule contains introns and corresponds to a DNA molecule in a generally one-to-one manner. An mRNA molecule corresponds to an HnRNA and DNA molecule from which the introns have been excised. A polynucleotide may consist of an entire gene, or any portion thereof. Operable anti-sense polynucleotides may comprise a fragment of the corresponding polynucleotide, and the definition of "polynucleotide" therefore includes all such operable anti-sense fragments.

The compositions and methods of the present invention also encompass variants of the above polypeptides and polynucleotides. A polypeptide "variant," as used herein, is a polypeptide that differs from the recited polypeptide only in conservative substitutions and/or modifications, such that the therapeutic, antigenic and/or immunogenic properties of the polypeptide are retained. In a preferred embodiment, variant polypeptides differ from an identified sequence by substitution, deletion or addition of five amino acids or fewer. Such variants may generally be identified by modifying one of the above polypeptide sequences, and evaluating the antigenic properties of the modified polypeptide using, for example, the representative procedures described herein. Polypeptide variants preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% identity (determined as described below) to the identified polypeptides.

As used herein, a "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. In general, the following groups of amino acids

represent conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his.

Variants may also, or alternatively, contain other modifications, including the deletion or addition of amino acids that have minimal influence on the antigenic properties, secondary structure and hydrophobic nature of the polypeptide. For example, a polypeptide may be conjugated to a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

A nucleotide "variant" is a sequence that differs from the recited nucleotide sequence in having one or more nucleotide deletions, substitutions or additions. Such modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983). Nucleotide variants may be naturally occurring allelic variants, or non-naturally occurring variants. Variant nucleotide sequences preferably exhibit at least about 70%, more preferably at least about 80% and most preferably at least about 90% identity (determined as described below) to the recited sequence.

The antigens provided by the present invention include variants that are encoded by DNA sequences which are substantially homologous to one or more of the DNA sequences specifically recited herein. "Substantial homology," as used herein, refers to DNA sequences that are capable of hybridizing under moderately stringent conditions. Suitable moderately stringent conditions include prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5X SSC, overnight or, in the event of cross-species homology, at 45°C with 0.5X SSC; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS. Such hybridizing DNA sequences are also within the scope of this invention, as are nucleotide sequences that, due to code degeneracy, encode an immunogenic polypeptide that is encoded by a hybridizing DNA sequence.

Two nucleotide or polypeptide sequences are said to be "identical" if the sequence of nucleotides or amino acid residues in the two sequences is the same when aligned for maximum correspondence as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, more preferably 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins - Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenesis pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) Fast and sensitive multiple sequence alignments on a microcomputer *CABIOS* 5:151-153; Myers, E.W. and Muller W. (1988) Optimal alignments in linear space *CABIOS* 4:11-17; Robinson, E.D. (1971) *Comb. Theor* 11:105; Santou, N. Nes, M. (1987) The neighbor joining method. A new method for reconstructing phylogenetic trees *Mol. Biol. Evol.* 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) *Numerical Taxonomy - the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) Rapid similarity searches of nucleic acid and protein data banks *Proc. Natl. Acad. Sci. USA* 80:726-730.

Preferably, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e. gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by

determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (i.e. the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Also included in the scope of the present invention are alleles of the genes encoding the nucleotide sequences recited herein. As used herein, an "allele" or "allelic sequence" is an alternative form of the gene which may result from at least one mutation in the nucleic acid sequence. Alleles may result in altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one, or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone or in combination with the others, one or more times in a given sequence.

For breast tumor polypeptides with immunoreactive properties, variants may, alternatively, be identified by modifying the amino acid sequence of one of the above polypeptides, and evaluating the immunoreactivity of the modified polypeptide. For breast tumor polypeptides useful for the generation of diagnostic binding agents, a variant may be identified by evaluating a modified polypeptide for the ability to generate antibodies that detect the presence or absence of breast cancer. Such modified sequences may be prepared and tested using, for example, the representative procedures described herein.

The breast tumor proteins of the present invention, and polynucleotide molecules encoding such proteins, may be isolated from breast tumor tissue using any of a variety of methods well known in the art. Polynucleotide sequences corresponding to a gene (or a portion thereof) encoding one of the inventive breast tumor proteins may be isolated from a breast tumor cDNA library using a subtraction technique as described in detail below. Examples of such DNA sequences are provided in SEQ ID NOS: 1- 94. Partial polynucleotide sequences thus obtained may be used to design oligonucleotide primers for the amplification of full-length polynucleotide sequences in a polymerase chain reaction (PCR), using techniques well known in the art (see, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY,

1989). Once a polynucleotide sequence encoding a polypeptide is obtained, any of the above modifications may be readily introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis as taught, for example, by Adelman et al. (*DNA*, 2:183, 1983).

The breast tumor polypeptides disclosed herein may also be generated by synthetic or recombinant means. Synthetic polypeptides having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may be generated using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain (see, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, CA), and may be operated according to the manufacturer's instructions.

Alternatively, any of the above polypeptides may be produced recombinantly by inserting a polynucleotide sequence that encodes the polypeptide into an expression vector and expressing the protein in an appropriate host. Any of a variety of expression vectors known to those of ordinary skill in the art may be employed to express recombinant polypeptides of this invention. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a polynucleotide molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line, such as CHO cells. The polynucleotide sequences expressed in this manner may encode naturally occurring polypeptides, portions of naturally occurring polypeptides, or other variants thereof.

In general, regardless of the method of preparation, the polypeptides disclosed herein are prepared in an isolated, substantially pure form (*i.e.*, the polypeptides are homogenous as determined by amino acid composition and primary sequence analysis). Preferably, the polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. In certain preferred embodiments,



described in more detail below, the substantially pure polypeptides are incorporated into pharmaceutical compositions or vaccines for use in one or more of the methods disclosed herein.

In a related aspect, the present invention provides fusion proteins comprising a first and a second inventive polypeptide or, alternatively, a polypeptide of the present invention and a known breast tumor antigen, together with variants of such fusion proteins.

A polynucleotide sequence encoding a fusion protein of the present invention is constructed using known recombinant DNA techniques to assemble separate polynucleotide sequences encoding the first and second polypeptides into an appropriate expression vector. The 3' end of a polynucleotide sequence encoding the first polypeptide is ligated, with or without a peptide linker, to the 5' end of a polynucleotide sequence encoding the second polypeptide so that the reading frames of the sequences are in phase to permit mRNA translation of the two DNA sequences into a single fusion protein that retains the biological activity of both the first and the second polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptides by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may be from 1 to about 50 amino acids in length. Peptide sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated polynucleotide sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of polynucleotides are located only 5' to the polynucleotide sequence encoding the first polypeptides. Similarly, stop codons require to end translation and transcription termination signals are only present 3' to the polynucleotide sequence encoding the second polypeptide.

Fusion proteins are also provided that comprise a polypeptide of the present invention together with an unrelated immunogenic protein. Preferably the immunogenic protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (see, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91 (1997)).

Polypeptides of the present invention that comprise an immunogenic portion of a breast tumor protein may generally be used for immunotherapy of breast cancer, wherein the polypeptide stimulates the patient's own immune response to breast tumor cells. In further aspects, the present invention provides methods for using one or more of the immunoreactive polypeptides encoded by a polynucleotide molecule having a sequence provided in SEQ ID NOS: 1- 94 (or fusion proteins comprising one or more such polypeptides and/or polynucleotides encoding such polypeptides) for immunotherapy of breast cancer in a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may be afflicted with a disease, or may be free of detectable disease. Accordingly, the above immunoreactive polypeptides (or fusion proteins or polynucleotide molecules encoding such polypeptides) may be used to treat breast cancer or to inhibit the development of breast cancer. The polypeptides may be administered either prior to or following surgical removal of primary tumors and/or treatment by administration of radiotherapy and conventional chemotherapeutic drugs.

In these aspects, the polypeptide or fusion protein is generally present within a pharmaceutical composition and/or a vaccine. Pharmaceutical compositions may comprise one or more polypeptides, each of which may contain one or more of the above sequences (or variants thereof), and a physiologically acceptable carrier. The vaccines may comprise one or more of such polypeptides and a non-specific immune response enhancer, wherein the non-

specific immune response enhancer is capable of eliciting or enhancing an immune response to an exogenous antigen. Examples of non-specific-immune response enhancers include adjuvants, biodegradable microspheres (e.g., polylactic galactide) and liposomes (into which the polypeptide is incorporated). Pharmaceutical compositions and vaccines may also contain other epitopes of breast tumor antigens, either incorporated into a combination polypeptide (i.e., a single polypeptide that contains multiple epitopes) or present within a separate polypeptide.

Alternatively, a pharmaceutical composition or vaccine may contain polynucleotides encoding one or more of the above polypeptides, such that the polypeptide is generated *in situ*. In such pharmaceutical compositions and vaccines, the polynucleotide may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid expression systems contain the necessary polynucleotide sequences for expression in the patient (such as a suitable promoter). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an epitope of a breast tumor cell antigen on its cell surface. In a preferred embodiment, the polynucleotide molecules may be introduced using a viral expression system (e.g., vaccinia or other pox virus, retrovirus, or adenovirus), which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N.Y. Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating polynucleotides into such expression systems are well known to those of ordinary skill in the art. The polynucleotides may also be "naked," as described, for example, in published PCT application WO 90/11092, and Ulmer et al., *Science* 259:1745-1749, 1993, reviewed by Cohen, *Science* 259:1691-1692,

1993. The uptake of naked polynucleotides may be increased by coating the polynucleotides onto biodegradable beads, which are efficiently transported into the cells.

Routes and frequency of administration, as well as dosage, will vary from individual to individual and may parallel those currently being used in immunotherapy of other diseases. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Between 1 and 10 doses may be administered over a 3-24 week period. Preferably, 4 doses are administered, at an interval of 3 months, and booster administrations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of polypeptide or polynucleotide molecule that is effective to raise an immune response (cellular and/or humoral) against breast tumor cells in a treated patient. A suitable immune response is at least 10-50% above the basal (*i.e.*, untreated) level. In general, the amount of polypeptide present in a dose (or produced *in situ* by the polynucleotide in a dose) ranges from about 1 pg to about 100 mg per kg of host, typically from about 10 pg to about 1 mg, and preferably from about 100 pg to about 1  $\mu$ g. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.01 mL to about 5 mL.

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a lipid, a wax and/or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and/or magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactic glycolide) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum

hydroxide or mineral oil, and a nonspecific stimulator of immune response, such as lipid A, *Bordella pertussis* or *Mycobacterium tuberculosis*. Such adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI) and Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ).

Polypeptides disclosed herein may also be employed in adoptive immunotherapy for the treatment of cancer. Adoptive immunotherapy may be broadly classified into either active or passive immunotherapy. In active immunotherapy, treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immune response-modifying agents (for example, tumor vaccines, bacterial adjuvants, and/or cytokines).

In passive immunotherapy, treatment involves the delivery of biologic reagents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (for example, CD8+ cytotoxic T-lymphocyte, CD4+ T-helper, tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells, lymphokine-activated killer cells), B cells, or antigen presenting cells (such as dendritic cells and macrophages) expressing the disclosed antigens. The polypeptides disclosed herein may also be used to generate antibodies or anti-idiotypic antibodies (as in U.S. Patent No. 4,918,164), for passive immunotherapy.

The predominant method of procuring adequate numbers of T-cells for adoptive immunotherapy is to grow immune T-cells *in vitro*. Culture conditions for expanding single antigen-specific T-cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. These *in vitro* culture conditions typically utilize intermittent stimulation with antigen, often in the presence of cytokines, such as IL-2, and non-dividing feeder cells. As noted above, the immunoreactive polypeptides described herein may be used to rapidly expand antigen-specific T cell cultures in order to generate sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage or B-cells, may be pulsed with immunoreactive polypeptides or transfected with a polynucleotide sequence(s), using standard techniques well known in the art. For example, antigen presenting cells may be transfected with a

polynucleotide sequence, wherein said sequence contains a promoter region appropriate for increasing expression, and can be expressed as part of a recombinant virus or other expression system. For cultured T-cells to be effective in therapy, the cultured T-cells must be able to grow and distribute widely and to survive long term *in vivo*. Studies have demonstrated that cultured T-cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (see, for example, Cheever, M., et al, "Therapy With Cultured T Cells: Principles Revisited," *Immunological Reviews*, 157:177, 1997).

The polypeptides disclosed herein may also be employed to generate and/or isolate tumor-reactive T-cells, which can then be administered to the patient. In one technique, antigen-specific T-cell lines may be generated by *in vivo* immunization with short peptides corresponding to immunogenic portions of the disclosed polypeptides. The resulting antigen specific CD8+ CTL clones may be isolated from the patient, expanded using standard tissue culture techniques, and returned to the patient.

Alternatively, peptides corresponding to immunogenic portions of the polypeptides may be employed to generate tumor reactive T cell subsets by selective *in vitro* stimulation and expansion of autologous T cells to provide antigen-specific T cells which may be subsequently transferred to the patient as described, for example, by Chang et al. (*Crit. Rev. Oncol. Hematol.*, 22(3), 213, 1996). Cells of the immune system, such as T cells, may be isolated from the peripheral blood of a patient, using a commercially available cell separation system, such as CellPro Incorporated's (Bothell, WA) CEPRATE™ system (see U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). The separated cells are stimulated with one or more of the immunoreactive polypeptides contained within a delivery vehicle, such as a microsphere, to provide antigen-specific T cells. The population of tumor antigen-specific T cells is then expanded using standard techniques and the cells are administered back to the patient.

In another embodiment, T-cell and/or antibody receptors specific for the polypeptides can be cloned, expanded, and transferred into other vectors or effector cells for use in adoptive immunotherapy.

In a further embodiment, syngeneic or autologous dendritic cells may be

pulsed with peptides corresponding to at least an immunogenic portion of a polypeptide disclosed herein. The resulting antigen-specific dendritic cells may either be transferred into a patient, or employed to stimulate T cells to provide antigen-specific T cells which may, in turn, be administered to a patient. The use of peptide-pulsed dendritic cells to generate antigen-specific T cells and the subsequent use of such antigen-specific T cells to eradicate tumors in a murine model has been demonstrated by Cheever et al, *Immunological Reviews*, 157:177, 1997).

Additionally, vectors expressing the disclosed polynucleotides may be introduced into stem cells taken from the patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

Polypeptides of the present invention may also, or alternatively, be used to generate binding agents, such as antibodies or fragments thereof, that are capable of detecting metastatic human breast tumors. Binding agents of the present invention may generally be prepared using methods known to those of ordinary skill in the art, including the representative procedures described herein. Binding agents are capable of differentiating between patients with and without breast cancer, using the representative assays described herein. In other words, antibodies or other binding agents raised against a breast tumor protein, or a suitable portion thereof, will generate a signal indicating the presence of primary or metastatic breast cancer in at least about 20% of patients afflicted with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without primary or metastatic breast cancer. Suitable portions of such breast tumor proteins are portions that are able to generate a binding agent that indicates the presence of primary or metastatic breast cancer in substantially all (*i.e.*, at least about 80%, and preferably at least about 90%) of the patients for which breast cancer would be indicated using the full length protein, and that indicate the absence of breast cancer in substantially all of those samples that would be negative when tested with full length protein. The representative assays described below, such as the two-antibody sandwich assay, may generally be employed for evaluating the ability of a binding agent to detect metastatic human breast tumors.

The ability of a polypeptide prepared as described herein to generate antibodies capable of detecting primary or metastatic human breast tumors may generally be evaluated by raising one or more antibodies against the polypeptide (using, for example, a representative method described herein) and determining the ability of such antibodies to detect such tumors in patients. This determination may be made by assaying biological samples from patients with and without primary or metastatic breast cancer for the presence of a polypeptide that binds to the generated antibodies. Such test assays may be performed, for example, using a representative procedure described below. Polypeptides that generate antibodies capable of detecting at least 20% of primary or metastatic breast tumors by such procedures are considered to be useful in assays for detecting primary or metastatic human breast tumors. Polypeptide specific antibodies may be used alone or in combination to improve sensitivity.

Polypeptides capable of detecting primary or metastatic human breast tumors may be used as markers for diagnosing breast cancer or for monitoring disease progression in patients. In one embodiment, breast cancer in a patient may be diagnosed by evaluating a biological sample obtained from the patient for the level of one or more of the above polypeptides, relative to a predetermined cut-off value. As used herein, suitable "biological samples" include blood, sera and urine.

The level of one or more of the above polypeptides may be evaluated using any binding agent specific for the polypeptide(s). A "binding agent," in the context of this invention, is any agent (such as a compound or a cell) that binds to a polypeptide as described above. As used herein, "binding" refers to a noncovalent association between two separate molecules (each of which may be free (*i.e.*, in solution) or present on the surface of a cell or a solid support), such that a "complex" is formed. Such a complex may be free or immobilized (either covalently or noncovalently) on a support material. The ability to bind may generally be evaluated by determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind" in the context of the present invention when the binding constant for complex formation



exceeds about  $10^3$  L/mol. The binding constant may be determined using methods well known to those of ordinary skill in the art.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome with or without a peptide component, an RNA molecule or a peptide. In a preferred embodiment, the binding partner is an antibody, or a fragment thereof. Such antibodies may be polyclonal, or monoclonal. In addition, the antibodies may be single chain, chimeric, CDR-grafted or humanized. Antibodies may be prepared by the methods described herein and by other methods well known to those of skill in the art.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding partner to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In a preferred embodiment, the assay involves the use of binding partner immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a second binding partner that contains a reporter group. Suitable second binding partners include antibodies that bind to the binding partner/polypeptide complex. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding partner after incubation of the binding partner with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding partner is indicative of the reactivity of the sample with the immobilized binding partner.

The solid support may be any material known to those of ordinary skill in the art to which the antigen may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as

adsorption, and covalent attachment (which may be a direct linkage between the antigen and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10  $\mu$ g, and preferably about 100 ng to about 1  $\mu$ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a second antibody (containing a reporter group) capable of binding to a different site on the polypeptide is added. The amount of second antibody that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior

to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is that period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include enzymes (such as horseradish peroxidase), substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. The conjugation of antibody to reporter group may be achieved using standard methods known to those of ordinary skill in the art.

The second antibody is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound second antibody is then removed and bound second antibody is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of breast cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the

cut-off value is the average mean signal obtained when the immobilized antibody is incubated with samples from patients without breast cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for breast cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (i.e., sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (i.e., the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for breast cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the antibody is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized antibody as the sample passes through the membrane. A second, labeled antibody then binds to the antibody-polypeptide complex as a solution containing the second antibody flows through the membrane. The detection of bound second antibody may then be performed as described above. In the strip test format, one end of the membrane to which antibody is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second antibody and to the area of immobilized antibody. Concentration of second antibody at the area of immobilized antibody indicates the presence of breast cancer. Typically, the concentration of second antibody at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of antibody immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody

sandwich assay, in the format discussed above. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1  $\mu$ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the antigens or antibodies of the present invention. The above descriptions are intended to be exemplary only.

In another embodiment, the above polypeptides may be used as markers for the progression of breast cancer. In this embodiment, assays as described above for the diagnosis of breast cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, breast cancer is progressing in those patients in whom the level of polypeptide detected by the binding agent increases over time. In contrast, breast cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

Antibodies for use in the above methods may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In one such technique, an immunogen comprising the antigenic polypeptide is initially injected into any of a wide variety of mammals (e.g., mice, rats, rabbits, sheep and goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for the antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.*

6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the polypeptide of interest). Such cell lines may be produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and tested for binding activity against the polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Monoclonal antibodies of the present invention may also be used as therapeutic reagents, to diminish or eliminate breast tumors. The antibodies may be used on their own (for instance, to inhibit metastases) or coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include  $^{90}\text{Y}$ ,  $^{123}\text{I}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{211}\text{At}$ , and  $^{212}\text{Bi}$ . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (e.g., covalently bonded) to a suitable monoclonal antibody either directly or indirectly (e.g., via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (e.g., a halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, e.g., U.S. Patent No. 4,671,958, to Rodwell et al.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (e.g., U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (e.g., U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (e.g., U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (e.g., U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed hydrolysis (e.g., U.S. Patent No. 4,569,789, to Blattler et al.).



It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Diagnostic reagents of the present invention may also comprise polynucleotide sequences encoding one or more of the above polypeptides, or one or more portions thereof. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify breast tumor-specific cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for a DNA molecule encoding a breast tumor protein of the present invention. The presence of the amplified



cDNA is then detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes specific for a DNA molecule encoding a breast tumor protein of the present invention may be used in a hybridization assay to detect the presence of an inventive polypeptide in a biological sample.

As used herein, the term "oligonucleotide primer/probe specific for a DNA molecule" means an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to the DNA molecule in question. Oligonucleotide primers and/or probes which may be usefully employed in the inventive diagnostic methods preferably have at least about 10-40 nucleotides. In a preferred embodiment, the oligonucleotide primers comprise at least about 10 contiguous nucleotides of a DNA molecule having a partial sequence selected from SEQ ID NOS: 1-94. Preferably, oligonucleotide probes for use in the inventive diagnostic methods comprise at least about 15 contiguous oligonucleotides of a DNA molecule having a partial sequence provided in SEQ ID NOS: 1-94. Techniques for both PCR based assays and hybridization assays are well known in the art (see, for example, Mullis *et al. Ibid*; Ehrlich, *Ibid*). Primers or probes may thus be used to detect breast tumor-specific sequences in biological samples, including blood, urine and/or breast tumor tissue.

The following Examples are offered by way of illustration and not by way of limitation.

## EXAMPLES

### Example 1

#### ISOLATION AND CHARACTERIZATION OF BREAST

#### TUMOR POLYPEPTIDES

This Example describes the isolation of breast tumor polypeptides from a breast tumor cDNA library.

A human breast tumor cDNA expression library was constructed from a pool of breast tumor poly A<sup>+</sup> RNA from three patients using a Superscript Plasmid System for

cDNA Synthesis and Plasmid Cloning kit (BRL Life Technologies, Gaithersburg, MD 20897) following the manufacturer's protocol. Specifically, breast tumor tissues were homogenized with polytron (Kinematica, Switzerland) and total RNA was extracted using Trizol reagent (BRL Life Technologies) as directed by the manufacturer. The poly A<sup>+</sup> RNA was then purified using a Qiagen oligotex spin column mRNA purification kit (Qiagen, Santa Clarita, CA 91355) according to the manufacturer's protocol. First-strand cDNA was synthesized using the NotI/Oligo-dT18 primer. Double-stranded cDNA was synthesized, ligated with EcoRI/BstX I adaptors (Invitrogen, Carlsbad, CA) and digested with NotI. Following size fractionation with Chroma Spin-1000 columns (Clontech, Palo Alto, CA 94303), the cDNA was ligated into the EcoRI/NotI site of pCDNA3.1 (Invitrogen, Carlsbad, CA) and transformed into ElectroMax *E. coli* DH10B cells (BRL Life Technologies) by electroporation.

Using the same procedure, a normal human breast cDNA expression library was prepared from a pool of four normal breast tissue specimens. The cDNA libraries were characterized by determining the number of independent colonies, the percentage of clones that carried insert, the average insert size and by sequence analysis. The breast tumor library contained  $1.14 \times 10^7$  independent colonies, with more than 90% of clones having a visible insert and the average insert size being 936 base pairs. The normal breast cDNA library contained  $6 \times 10^6$  independent colonies, with 83% of clones having inserts and the average insert size being 1015 base pairs. Sequencing analysis showed both libraries to contain good complex cDNA clones that were synthesized from mRNA, with minimal rRNA and mitochondrial DNA contamination sequencing.

cDNA library subtraction was performed using the above breast tumor and normal breast cDNA libraries, as described by Hara *et al.* (*Blood*, 84:189-199, 1994) with some modifications. Specifically, a breast tumor-specific subtracted cDNA library was generated as follows. Normal breast cDNA library (70  $\mu$ g) was digested with EcoRI, NotI, and SfiI, followed by a filling-in reaction with DNA polymerase Klenow fragment. After phenol-chloroform extraction and ethanol precipitation, the DNA was dissolved in 100  $\mu$ l of H<sub>2</sub>O, heat-denatured and mixed with 100  $\mu$ l (100  $\mu$ g) of Photoprobe biotin (Vector Laboratories, Burlingame, CA), the resulting mixture was irradiated with a 270 W sunlamp

on ice for 20 minutes. Additional Photoprobe biotin (50  $\mu$ l) was added and the biotinylation reaction was repeated. After extraction with butanol five times, the DNA was ethanol-precipitated and dissolved in 23  $\mu$ l H<sub>2</sub>O to form the driver DNA.

To form the tracer DNA, 10  $\mu$ g breast tumor cDNA library was digested with BamHI and XhoI, phenol chloroform extracted and passed through Chroma spin-400 columns (Clontech). Following ethanol precipitation, the tracer DNA was dissolved in 5  $\mu$ l H<sub>2</sub>O. Tracer DNA was mixed with 15  $\mu$ l driver DNA and 20  $\mu$ l of 2 x hybridization buffer (1.5 M NaCl/10 mM EDTA/50 mM HEPES pH 7.5/0.2% sodium dodecyl sulfate), overlaid with mineral oil, and heat-denatured completely. The sample was immediately transferred into a 68 °C water bath and incubated for 20 hours (long hybridization [LH]). The reaction mixture was then subjected to a streptavidin treatment followed by phenol/chloroform extraction. This process was repeated three more times. Subtracted DNA was precipitated, dissolved in 12  $\mu$ l H<sub>2</sub>O, mixed with 8  $\mu$ l driver DNA and 20  $\mu$ l of 2 x hybridization buffer, and subjected to a hybridization at 68 °C for 2 hours (short hybridization [SH]). After removal of biotinylated double-stranded DNA, subtracted cDNA was ligated into BamHI/XhoI site of chloramphenicol resistant pBCSK<sup>+</sup> (Stratagene, La Jolla, CA 92037) and transformed into ElectroMax *E. coli* DH10B cells by electroporation to generate a breast tumor specific subtracted cDNA library.

To analyze the subtracted cDNA library, plasmid DNA was prepared from 100 independent clones, randomly picked from the subtracted breast tumor specific library and characterized by DNA sequencing with a Perkin Elmer/Applied Biosystems Division Automated Sequencer Model 373A (Foster City, CA). Thirty-eight distinct cDNA clones were found in the subtracted breast tumor-specific cDNA library. The determined 3' cDNA sequences for 14 of these clones are provided in SEQ ID NO: 1-14, with the corresponding 5' cDNA sequences being provided in SEQ ID NO: 15-28, respectively. The determined one strand (5' or 3') cDNA sequences for the remaining clones are provided in SEQ ID NO: 29-52. Comparison of these cDNA sequences with known sequences in the gene bank using the EMBL and GenBank databases (Release 97) revealed no significant homologies to the sequences provided in SEQ ID NO: 3, 10, 17, 24 and 45-52. The sequences provided in SEQ ID NO: 1, 2, 4-9, 11-16, 18-23, 25-41, 43 and 44 were found to show at least some degree of

homology to known human genes. The sequence of SEQ ID NO: 42 was found to show some homology to a known yeast gene.

Data was analyzed using Synteni provided GEMTOOLS Software. Twenty one distinct cDNA clones were found to be over-expressed in breast tumor and expressed at low levels in all normal tissues tested. The determined partial cDNA sequences for these clones are provided in SEQ ID NO: 53- 73. Comparison of the sequences of SEQ ID NO: 53, 54, and 68-71 with those in the gene bank as described above, revealed some homology to previously identified human genes. No significant homologies were found to the sequences of SEQ ID NO: 55-67, 72 (referred to as JJ 9434,7117), and 73 (referred to as B535S).

In a further experiment, cDNA fragments analyzed by DNA microarray were obtained from two subtraction libraries derived by conventional subtraction, as described above. In one instance the tester was derived from primary breast tumors. In the second instance, a metastatic breast tumor was employed as the tester. Drivers consisted of normal breast.

cDNA fragments from these two libraries were submitted as templates for DNA microarray analysis. DNA chips were analyzed by hybridizing with fluorescent probes derived from mRNA from both tumor and normal tissues. Analysis of the data was accomplished by creating three groups from the sets of probes. The composition of these probe groups, referred to as Breast Tumor/mets, Normal non-breast tissues, and Metastatic breast tumors. Two comparisons were performed using the modified Gemtools analysis. The first comparison was to identify templates with elevated expression in breast tumors. The second was to identify templates not recovered in the first comparison that yielded elevated expression in metastatic breast tumors. An arbitrary level of increased expression (mean of tumor expression versus the mean of normal tissue expression) was set at approximately 2.2.

In the first round of comparison to identify overexpression in breast tumors, two novel gene sequences were identified, hereinafter referred to as B534S and B538S (SEQ ID NO: 89 and 90), and six sequences that showed some degree of homology to previously identified genes (SEQ ID NO: 74-79). Additionally, in a second comparison to identify elevated expression in metastatic breast tumors, five novel sequences were identified, hereinafter referred to as B535S (overexpressed in this analysis as well as what was described

above), B542S, B543S, P501S, and B541S (SEQ ID NO: 73, and 91-94), as well as nine gene sequences that showed some homology to known genes (SEQ ID NO: 80-88). Clone B534S and B538S (SEQ ID NO: 89 and 90) were shown to be overexpressed in both breast tumors and metastatic breast tumors.

### Example 2

#### GENERATION OF HUMAN CD8+ CYTOTOXIC T-CELLS THAT RECOGNIZE ANTIGEN PRESENTING CELLS EXPRESSING BREAST TUMOR ANTIGENS

This Example illustrates the generation of T cells that recognize target cells expressing the antigen B511S, also known as 1016-F8 (SEQ ID NO: 56). Human CD8+ T cells were primed *in-vitro* to the B511S gene product using dendritic cells infected with a recombinant vaccinia virus engineered to express B511S as follows (also see Yee et al., Journal of Immunology (1996) 157 (9):4079-86). Dendritic cells (DC) were generated from peripheral blood derived monocytes by differentiation for 5 days in the presence of 50 µg/ml GM-CSF and 30 µg/ml IL-4. DC were harvested, plated in wells of a 24-well plate at a density of  $2 \times 10^5$  cells/well and infected for 12 hours with B511S expressing vaccinia at a multiplicity of infection of 5. DC were then matured overnight by the addition of 3 µg/ml CD40-Ligand and UV irradiated at 100µW for 10 minutes. CD8+ T cells were isolated using magnetic beads, and priming cultures were initiated in individual wells (typically in 24 wells of a 24-well plate) using  $7 \times 10^5$  CD8+ T cells and  $1 \times 10^6$  irradiated CD8-depleted PBMC; IL-7 at 10 ng/ml was added to cultures at day 1. Cultures were re-stimulated every 7-10 days using autologous primary fibroblasts retrovirally transduced with B511S and the costimulatory molecule B7.1. Cultures were supplemented at day 1 with 15 I.U. of IL-2. Following 4 such stimulation cycles, CD8+ cultures were tested for their ability to specifically recognize autologous fibroblasts transduced with B511S using an interferon-γ Elispot assay (see Lalvani et al J. Experimental Medicine (1997) 186:859-965). Briefly, T cells from individual microcultures were added to 96-well Elispot plates that contained autologous fibroblasts transduced to express either B511S or as a negative control antigen EGFP, and incubated overnight at 37° C; wells also contained IL-12 at 10 ng/ml. Cultures were identified that specifically produced interferon-γ only in response to B511S transduced

fibroblasts; such lines were further expanded and also cloned by limiting dilution on autologous B-LCL retrovirally transduced with B511S. Lines and clones were identified that could specifically recognize autologous B-LCL transduced with B511S but not autologous B-LCL transduced with the control antigens EGFP or HLA-A3. An example demonstrating the ability of human CTL cell lines derived from such experiments to specifically recognize and lyse B511S expressing targets is presented in Figure 1.

### Example 3

#### SYNTHESIS OF POLYPEPTIDES

Polypeptides may be synthesized on an Perkin Elmer/Applied Biosystems Division 430A peptide synthesizer using Fmoc chemistry with HPTU (O-Benzotriazole-N,N,N',N'-tetramethyluronium hexafluorophosphate) activation. A Gly-Cys-Gly sequence may be attached to the amino terminus of the peptide to provide a method of conjugation, binding to an immobilized surface, or labeling of the peptide. Cleavage of the peptides from the solid support may be carried out using the following cleavage mixture: trifluoroacetic acid:ethanedithiol:thioanisole:water:phenol (40:1:2:2:3). After cleaving for 2 hours, the peptides may be precipitated in cold methyl-t-butyl-ether. The peptide pellets may then be dissolved in water containing 0.1% trifluoroacetic acid (TFA) and lyophilized prior to purification by C18 reverse phase HPLC. A gradient of 0%-60% acetonitrile (containing 0.1% TFA) in water (containing 0.1% TFA) may be used to elute the peptides. Following lyophilization of the pure fractions, the peptides may be characterized using electrospray or other types of mass spectrometry and by amino acid analysis.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

## CLAIMS

1. An isolated polypeptide comprising an immunogenic portion of a breast protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, and 89-94; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

2. An isolated polynucleotide molecule comprising a nucleotide sequence encoding the polypeptide of claim 1.

3. An isolated polynucleotide molecule comprising a sequence provided in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, and 89-94.

4. An expression vector comprising a polynucleotide molecule according to any one of claims 2 and 3.

5. A host cell transformed with the expression vector of claim 4.

6. The host cell of claim 5 wherein the host cell is selected from the group consisting of *E. coli*, yeast and mammalian cell lines.

7. A pharmaceutical composition comprising the polypeptide of claim 1 and a physiologically acceptable carrier.

8. A vaccine comprising the polypeptide of claim 1 and a non-specific immune response enhancer.

9. The vaccine of claim 8 wherein the non-specific immune response enhancer is an adjuvant.

10. A vaccine comprising a polynucleotide molecule of any one of claims 2 and 3 and a non-specific immune response enhancer.

11. The vaccine of claim 10 wherein the non-specific immune response enhancer is an adjuvant.

12. A pharmaceutical composition for the treatment of breast cancer comprising a polypeptide and a physiologically acceptable carrier, the polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

13. A vaccine for the treatment of breast cancer comprising a polypeptide and a non-specific immune response enhancer, said polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

14. The vaccine of claim 13 wherein the non-specific immune response enhancer is an adjuvant.

15. A vaccine for the treatment of breast cancer comprising a polynucleotide molecule and a non-specific immune response enhancer, the polynucleotide



molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

16. The vaccine of claim 15, wherein the non-specific immune response enhancer is an adjuvant.

17. A pharmaceutical composition according to claims 7 or 12, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

18. A vaccine according to any one of claims 8, 10, 13 or 15, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

19. A fusion protein comprising at least one polypeptide according to claim 1.

20. A pharmaceutical composition comprising a fusion protein according to claim 19 and a physiologically acceptable carrier.

21. A vaccine comprising a fusion protein according to claim 19 and a non-specific immune response enhancer.

22. The vaccine of claim 21 wherein the non-specific immune response enhancer is an adjuvant.

23. A pharmaceutical composition according to claim 20, for use in manufacture of a medicament for inhibiting the development of breast cancer in a patient.

24. A vaccine according to claim 21, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

25. A method for detecting breast cancer in a patient, comprising:

(a) contacting a biological sample from a patient with a binding agent which is capable of binding to a polypeptide, the polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence provided in SEQ ID NO: 1-94 under moderately stringent conditions; and

(b) detecting in the sample a protein or polypeptide that binds to the binding agent, thereby detecting breast cancer in the patient.

26. The method of claim 25 wherein the binding agent is a monoclonal antibody.

27. The method of claim 26 wherein the binding agent is a polyclonal antibody.

28. A method for monitoring the progression of breast cancer in a patient, comprising:

(a) contacting a biological sample from a patient with a binding agent that is capable of binding to a polypeptide, said polypeptide comprising an immunogenic portion of a breast protein, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence provided in SEQ ID NO: 1-94 under moderately stringent conditions;

(b) determining in the sample an amount of a protein or polypeptide that binds to the binding agent;

(c) repeating steps (a) and (b); and

(d) comparing the amount of polypeptide detected in steps (b) and (c) to monitor the progression of breast cancer in the patient.

29. A monoclonal antibody that binds to a polypeptide comprising an immunogenic portion of a breast protein or a variant of said protein that differs only in conservative substitutions and/or modifications, wherein said protein comprises an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of: (a) nucleotide sequences recited in SEQ ID NOS: 3, 10, 17, 24, 45-52, 55-67, 72, 73, and 89-94; (b) complements of said nucleotide sequences; and (c) sequences that hybridize to a sequence of (a) or (b) under moderately stringent conditions.

30. A monoclonal antibody according to claim 29, for use in the manufacture of a medicament for inhibiting the development of breast cancer in a patient.

31. The monoclonal antibody of claim 30 wherein the monoclonal antibody is conjugated to a therapeutic agent.

32. A method for detecting breast cancer in a patient comprising:

(a) contacting a biological sample from a patient with at least two oligonucleotide primers in a polymerase chain reaction, wherein at least one of the oligonucleotides is specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NO: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions; and

(b) detecting in the sample a polynucleotide sequence that amplifies in the presence of the oligonucleotide primers, thereby detecting breast cancer.

33. The method of claim 32, wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from SEQ ID NOS: 1-94.

34. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies of claim 29; and
- (b) a detection reagent.

35. A diagnostic kit comprising:

- (a) one or more monoclonal antibodies that bind to a polypeptide encoded by a polynucleotide molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, and 74-88, complements of said sequences and sequences that hybridize to a sequence of SEQ ID NO: 1, 2, 4-9, 11-16, 18-23, 25-44, 53, 54, 68-71, or 74-88 under moderately stringent conditions; and
- (b) a detection reagent.

36. The kit of claims 34 or 35 wherein the monoclonal antibodies are immobilized on a solid support.

37. The kit of claim 36 wherein the solid support comprises nitrocellulose, latex or a plastic material.

38. The kit of claims 34 or 35 wherein the detection reagent comprises a reporter group conjugated to a binding agent.

39. The kit of claim 38 wherein the binding agent is selected from the group consisting of anti-immunoglobulins, Protein G, Protein A and lectins.

40. The kit of claim 38 wherein the reporter group is selected from the group consisting of radioisotopes, fluorescent groups, luminescent groups, enzymes, biotin and dye particles.

41. A diagnostic kit comprising at least two oligonucleotide primers, at least one of the oligonucleotide primers being specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein

comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions.

42. A diagnostic kit of claim 41 wherein at least one of the oligonucleotide primers comprises at least about 10 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from SEQ ID NOS: 1-94.

43. A method for detecting breast cancer in a patient, comprising:

- (a) obtaining a biological sample from the patient;
- (b) contacting the biological sample with an oligonucleotide probe specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions; and
- (c) detecting in the sample a polynucleotide sequence that hybridizes to the oligonucleotide probe, thereby detecting breast cancer in the patient.

44. The method of claim 43 wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from the group consisting of SEQ ID NOS: 1-94.

45. A diagnostic kit comprising an oligonucleotide probe specific for a polynucleotide molecule encoding a polypeptide comprising an immunogenic portion of a breast protein, said protein comprising an amino acid sequence encoded by a polynucleotide molecule comprising a sequence selected from the group consisting of nucleotide sequences recited in SEQ ID NOS: 1-94, complements of said nucleotide sequences, and sequences that hybridize to a sequence of SEQ ID NO: 1-94 under moderately stringent conditions.

46. The diagnostic kit of claim 45, wherein the oligonucleotide probe comprises at least about 15 contiguous nucleotides of a polynucleotide molecule comprising a sequence selected from the group consisting of SEQ ID NOS: 1-94.

47. Peripheral blood cells from a patient incubated in the presence of at least one polypeptide of claim 1, such that T cells proliferate, for use in the manufacture of a medicament for treating breast cancer in a patient.

48. The blood cells of claim 47 wherein the T cells is repeated one or more times.



49. A composition for the treatment of breast cancer in a patient, comprising T cells proliferated in the presence of a polypeptide of claim 1, in combination with a pharmaceutically acceptable carrier.

50. An antigen presenting cells incubated in the presence of at least one polypeptide of claim 1, for use in the manufacture of a medicament for treating breast cancer in a patient.

51. The cells of claim 50 wherein the antigen presenting cells are selected from the group consisting of dendritic and macrophage cells.

52. A composition for the treatment of breast cancer in a patient, comprising antigen presenting cells incubated in the presence of a polypeptide of claim 1, in combination with a pharmaceutically acceptable carrier.

1/1

*Fig. 1A**Fig. 1B*

## SEQUENCE LISTING

&lt;110&gt; Corixa Corporation

&lt;120&gt; COMPOUNDS FOR IMMUNOTHERAPY AND DIAGNOSIS OF BREAST CANCER AND METHODS FOR THEIR USE

&lt;130&gt; 210121.446PC

&lt;140&gt; PCT

&lt;141&gt; 1998-12-22

&lt;160&gt; 94

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 402

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 1

```
tttttttttt tttttaggag aactgaatca aacagatttt attcaacttt ttagatgagg 60
aaaacaaatn atacgaaatn ngtcataaga aatgctttct tataaccacta tctcaaacca 120
ctttcaatat tttaaaaaat gctcacgcag caaatatgaa aagctncaac acttcccttt 180
gttaacttgc tgcaatnaat gcaactttaa canacatata aatttcttct gtatcttaaa 240
agttnaatta ctaattttta tgatnttnt caagatnttt attcatatac tttaaatgac 300
tcnttgccna tacatacnta ttttctttac ttttttttta cnatnggcc aacagctttca 360
ngcagncnc aaaaatctta ccggttaatt acacggggtt gt 402
```

&lt;210&gt; 2

&lt;211&gt; 424

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```
tttttttttt ttttttaag gtacacattt ctttttcatt ctgttttnatg cagcaaataa 60
ttcgtttggca tcttctctgt gatgggcagc ttgctaaaat taactcagg ccccttagct 120
ncatttccaa ctgagccac gctttcaacc nngccnaaca aagaaaatca gttngggtta 180
aattctttgc tgganacaaa gaactacatt cttttgtaaa tnatgctttg tttgctctgt 240
gcaaacncag attgaaggga anaagganac ttntggggac ggaaacaact ngnagaagca 300
gganccgcc agggncattt cctcaccatg cttaatcttg cncctacttg cngggcacca 360
ttaaacttgg tgcaaaaggc gcaattggtg nanggaaccc cacaccttcc ttaaaaagca 420
gggc 424
```

&lt;210&gt; 3

&lt;211&gt; 421

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 3

```
tttttttttt tttttccaa tttaaaaag cttttttcat acttcaatta caccanactt 60
aatnatttca tgagtaaact ngacattatt atttnaaaat ttgcatattt aaaatttgn 120
tcanttactt ccagactgtt tgcanaatga agggaggatc actcaagngc tgatctenca 180
ctntctgcag tctnctgtcc tgtgcccgn ctaatggatc gacactanat ggacagntcn 240
```



cagatcttcc gttcttntcc tttccccaat ttncaccnc tccccttctt ncccggatcn 300  
tttggggaca tgntaatttt gcnatcctta aaccctgcc gccangggc ccanctcag 360  
gggtgggttaa tgttcgneng gcttnttgac cncctgcc ctttnantcc naacccaag 420  
c 421

&lt;210&gt; 4

&lt;211&gt; 423

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 4

tttttttatt tttttttcta tttntntat ttntgnggt tctgtgtgt aattagnang 60  
tgtgtatgcg tangtaenta tgtntgcata tttaacctgt tncctttcca tttttaaaat 120  
aaaatctcaa natngtantt ggttnatggg agtaaanaga gactatngat naattttaac 180  
atggacacng tgaaatgtag ccgctnatca nttaaaaact tcaatttgaa ggcttttnc 240  
cctccnaata aaaatnccng gccctactgg gttaagcaac attgcatntc taaagaaacc 300  
acatgcanac nagttaaac tgtnactgg tcangcaaac cnaantggaa nanaagggnn 360  
ttnccccan ggacantcng aatttttta acaaattacn atncccccc ngggggagcc 420  
tgt 423

&lt;210&gt; 5

&lt;211&gt; 355

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

acgaccacct natttcgtat ctttcaactc ttttcgaccg gacctcttat tcggaagcgt 60  
tccaggaaga caggtctcaa cttagggatc agatcacgtt atcaacgctc tgggatcgct 120  
gcaacctggc acttcaagga agtgcaccga tnacgtctag accggccaac acagatctag 180  
aggtggccaa ctgatcactg taggagctga ctggcaanan tcaaccgggc cccaaccnag 240  
agtgaccaan acnaccattn aggatcacc acaggcactc ctctcctag ggccaaccna 300  
ccaaacggct ggccaatggg ggggtttaat atttggttna aaaattgatt ttaaa 355

&lt;210&gt; 6

&lt;211&gt; 423

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 6

tttttttttt ttttttgaca ggaagtaaaa ttattggtn antattaana ggggggcagc 60  
acattggaag cctcatgan tgcagggccc gccacttgct cagagggcca cnattggga 120  
tgtacttaac cccacagecn tctgggatna gccgcttttc agccaccatn ttttcaaatt 180  
catcagcatt aaacttggtt aanncccaact tctttaagat ntgnatcttc tggcgccag 240  
naaacttgaa ctggccttg cgcagggcct caatcacatg ctcttggttc tgcagcttg 300  
tgcgnaagga cntaactnact tggccnatgt gaacctggc cacantgcc tggggcttc 360  
caaagcacc tcgcaagcct ntttggance tgnccgcccc ngcacaggga caacatcttg 420  
ttt 423

&lt;210&gt; 7

&lt;211&gt; 410

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 7

ttcgactgg ctaaaacaaa ccgccttgca agttngaaa aatttatcaa tggaccaaat 60

aatgctcata tccnacaagt tggtagcctg tnttatnata aaaaaatgta tnatgctcct 120  
nanttggtgt acaataatgt tccaatttng gacnttcggc atctaccctg gttcacctgg 180  
gtaaatatca ggcagctttt gatggggcta ggaaagctaa cagtactcga acatgggaaa 240  
gaggtctgct tcgccngtgt anatgggaaa naattccgtc ttgctcngat ttgtggactt 300  
catattgttg tacatgcaga tgaatnngaa gaacttgta actactatca ggatcgtggc 360  
tttttnnaaa agctnatcac catgttgga ggggactng gacttgagcg 410

&lt;210&gt; 8

&lt;211&gt; 274

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

tttttttttt ttttaggtc atacatattt tttattataa canatatntg tatatacata 60  
taatatatgt gtatatatcc acgtgtgtgt gtgtgtatca aaaaacaacan aanttttagtg 120  
atctatatct ntngctcaca tatgcatggg agataccagt aaaaaataag tnaatctcca 180  
taatatgttt taaaactcan anaaatcnga gagactnaaa gaaaacgtnn atcannatga 240  
ttgtngataa tcttgaanaa tnacnaaaac atat 274

&lt;210&gt; 9

&lt;211&gt; 322

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

tttttttttt ttttgtgctt tattgcaccg gcnaaaactt ctgacactat attaaactca 60  
ataagagtga taagtgtgaa aatccttgcc ttctctttaa tcttaatgna naggcatctg 120  
gtttttcacc attaanatga ataattggctn tatgtatttt tatnnatggg cttnatggag 180  
ttaaaaaagt ttctctctnt cctngttat ctaanagttt tnatcaaaaa tgggtataat 240  
attngttca gtacttttnc ctgcacctat agatatgatn ctgttatttt ttctcttng 300  
cctnnanata tgatggatna ca 322

&lt;210&gt; 10

&lt;211&gt; 425

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 10

tttttttttt tttttattct gcagccatta aatgctgaac actagatnct tttttgtgga 60  
ggtcacaaaa taagtacaga atatnacaca cgccctgccc ataaaaagca cagctcccag 120  
ttctataattt acaatatctc tgggaattcca ccttccttc taatttgact aatatttctg 180  
cttctcaggc agcagcgctt tctggcaacc ataagaacca acntgnggac taggtcggtg 240  
ggccaaggat caggaaacag aanaatggaa gnagcccccn tgacnctatt aancntnaa 300  
actatctnaa ctgctagttt tcaggcttta aatcatgtaa natacgtgtc cttnttgctg 360  
caaccggaag catcctagat ggtacactct ctccagggtc caggaaaaga tcccaaatng 420  
caggn 425

&lt;210&gt; 11

&lt;211&gt; 424

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

ttttnttant ttttttance nctnntccnn tntgttgnag ggggtaccaa atttctttat 60  
ttaaaggaat ggtacaaatc aaaaaactta atttaatttt tnggtacaac ttatagaaaa 120

ggttaaggaa accccaacat gcatgcactg ccttggtaac cagggnattc ccccneggt 180  
ntggggaaat tagcccaang ctnagctttc attatcactn tccccaggg tntgcttttc 240  
aaaaaaattt nccgccnagc cnaatccggg cncctccatc tggcgcaant tggcacttg 300  
gtccccnat tctttaangg cttncacctn ctcatcggg tnatgtgtct caattaaatc 360  
ccacngatgg gggtcatttt tntcnnttag ccagtttggt nagttccgtt attganaaaa 420  
ccan 424

&lt;210&gt; 12

&lt;211&gt; 426

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

tttttttttt ttttnttaaa aagcttttat ctctgctta cattacccat ctgttcttgc 60  
atgttgctcg cttttccac tagagccctt aacaacttaa tcatggttat ttttaagggt 120  
ctaataattc cnaaactggg atcataaata agtctcggtc tnatgcttgc tttctctcta 180  
tcacactgtg ttngttgctt tttnacatgc tttgtaattt ttggtgaaa gctgaaaaat 240  
nacatacctg gttntacaac ctgaggtaan cagccttnta gtgtgaggtt ttatatntta 300  
ctggctaaga gctnggcncct gttnantant tgttgtaant ntatatgcca naggtctttna 360  
ttccnctng tgtccttgc tnatgtaccc attnttttag gggttcccta naaactctat 420  
ctnaat 426

&lt;210&gt; 13

&lt;211&gt; 419

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 13

tttttttttt ttttnagat agactctcac tctttgccc aggctggagt gcagtggcgc 60  
aatcaaggct cactgcaacc tctgccttat aaagcatttn cttaaaggtaac aagctaaatt 120  
ttaaaaaat ctctncacaa ctaatgtata acaaaaatta gttctacctc ataaacnct 180  
ggctcagccc tcgnaacaca tttccctggt ctcaactgat gaacactcca naaacagAAC 240  
anatntaagc tttccaggc ccagaaaagc tcgcgagggg atttgctntg tgtgtgacac 300  
acttgccacc ctgtgagcgc acagctccac acntgctttg ggccgcattt gcaagttctc 360  
tgtaancccc ctgnaagacc cggatcagct gggtnгааat tgcangcnc tttttggca 419

&lt;210&gt; 14

&lt;211&gt; 400

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 14

aanccattgc caagggtatc cggaggattg tggctgtcac aggtncggag gcccahaagg 60  
ccctcaggaa agcaaaagagc ttgaaaaatg tctctctgtc atggaagccn aagtgaaggc 120  
tcanactgct ccaacaaggga tntgcanagg gagatcgcta accttgaggga ggccctgggc 180  
actgcagtcn tccccantg gcagaaggat gaattgcggg agactctcan atcccttang 240  
gaaggtcgtg gatnacttg accgagcctc nnaagccaat ntccagaaca agtgttgag 300  
aagacaaagc anttcatcga cgccaacccc naccggcctc tnttctctg ganattgana 360  
gcggcgcccc cgccagggc cttaataanc cntgaagctn 400

&lt;210&gt; 15

&lt;211&gt; 395

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

<400> 15  
 tgctttgctg cgtccaggaa gattagatng aanaatacat attgatttgc caaatgaaca 60  
 agcgagatta gaentactga anateccatgc aggtccatt acaaagcatg gtgaaataga 120  
 tgatgaagca attgtgaagc tatcggatgg ctttnatgga gcagatctga gaaatgtttg 180  
 tactgaagca ggtatgttcg caattcgtgc tgatcatgat tttgtagtac aggaagactt 240  
 catgaaagcn gtcagaanag tggctnattc tnaaagctgg agtctaaatt ggacnacnac 300  
 ctntgtatct actgttggan ttttgatgct gcatgacaga ttttgcttan tgtaaaaatn 360  
 aagttcaaga aaattatggt agttttggcc attat 395

<210> 16

<211> 404

<212> DNA

<213> Homo sapiens

<400> 16  
 ccaccactaa aatectggct gagccctacn agtacctgtg cccctccccc aggacgagat 60  
 nagggcacac cctttaagtn aggtgacagg tcacctttaa gtgaggacag tcagctnaat 120  
 ttcacctett gggcttgagt acctggttct cgtgccctga ggcgacnctn agccctgcag 180  
 ctncatgta cgtgctgcca atngtcttga tcttctccac gccnctnaac ttgggcttca 240  
 gtaggagctg caggcnagaa ngaagcgggt aacagcgcca ctccatagcc gcagccnggc 300  
 tgcccctgct tctcaaggag ggggtgtggg ttcctccacc atcgccgcc ttgcaaacac 360  
 ntctcanggc ttcctnccg gctnancgca ngacttaagc atgg 404

<210> 17

<211> 360

<212> DNA

<213> Homo sapiens

<400> 17  
 ggccagaagc tttccacaaa ccagtgaagg tggcagcaaa gaaagcctct tagacnagga 60  
 gctggcagca gctgctatct ngatngacng cagaaccaa ccactaatc agcaaacaca 120  
 acctcatacc tnaccgcttc ccttnaatg gccttcggtg tgtgcgca tgggcacgtg 180  
 cggggagaac catacttatt cccctnttcc cggcctacca cctctnctcc cccttctctt 240  
 ctctncaatt actntctccn ctgcttntt ctanaccta ctgctngtnt cnanagccng 300  
 cccgcaatta cctggcaaaa ctgcgcaccc ttcgggcagc gctaaanaat gcacatttac 360

<210> 18

<211> 316

<212> DNA

<213> Homo sapiens

<400> 18  
 atacatatac acatatatga ttttagatag agccatatac ctngaagtag tanatttgtt 60  
 tgtgtgtata tgtatgtgc tactcatttt aaataaactt gtgatagaga tgtaattntg 120  
 agccagtttt tcatttgctt aaatnactca ccaagtaact aattaagtn tctttactct 180  
 taatgtnnag tagtgagatt ctgttgaagg tgatattaaa aaccattcta tattaattaa 240  
 cattcatggt gtttttataa agcttatttg aaatcnaatt atgattatct ttcataccag 300  
 tcgatnttat gtangt 316

<210> 19

<211> 350

<212> DNA

<213> Homo sapiens

<400> 19

aagggatgca nataatgctg tgtatgagct tgatggaaaa gaactctgta gtgaaagggg 60  
tactattgaa catgctnngg ctcggtcacg aggtggaaaga ggttagaggac gatactctga 120  
ccggttttagt agtcgcagac ctcgaaatga tagacgaaat gctccacctg taagaacaga 180  
anacgtctt atagttgaga atttatcttc aagagtcagc tggcagggtt gttganatac 240  
agttttgagt tnttttgatg tggcttttta aaaaagttat gggttactna tggtatattg 300  
ttttattaaa agtagttttn aattaatgga tntgatggaa ttgttggttt 350

&lt;210&gt; 20

&lt;211&gt; 367

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 20

gntnnnnhca agatcctnct ntccccngg gngcccccnc cncngtnat naccgggttt 60  
ntaanatcnn gccgnccecg aagtctcnc nntgccgaga tgncccttat ncnncnnatgn 120  
ncaattntga cctnnggcga anaatggcng nngtgtatca gtntccnctc tngggntctt 180  
tagnatctga ccactangac cncctatcct ctcaaacctt gtanncngcc ctaatttggtg 240  
ccaattagtg catgntanag cntcctggcc cagatggcct ccatatcctg gtncggcttc 300  
cgccccctacc angncatccn catctactag agcttatccg ctncntgngg cgcaccggnt 360  
ccccnct 367

&lt;210&gt; 21

&lt;211&gt; 366

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 21

cccaacacaa tggcttaagt anaactgtat tgctctgtag tatagttcca cattggcaac 60  
ctacaatggg aaaatccata cataagtcag ttacttctcn atgagctttc tccttctgaa 120  
tcctttatct tctgaagaaa gtacacacct tggtnatgat atctttgaat tgccttctt 180  
tccaggcatc agttggatga ttcacatggt taattatggc attatcatat tcttcatact 240  
tgtcatatga aaacaccagt tctgccnna gatgagcttg ttctgcagct cttagcacct 300  
tggaatatt cactctagac cagaaacagc tcccggtgct ccctcatttt ctgaggctta 360  
aatttn 366

&lt;210&gt; 22

&lt;211&gt; 315

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 22

acttaatgca atctctggag gataatttgg atcaagaaat aaagaanaaa tgaattagga 60  
gaagaaatna ctgggtnata tttcaatatt ttagaacttt aanaatgttg actatgattt 120  
caatatattt gtnaaaactg agatacangt ttgacctata tctgcatttt gataattaaa 180  
cnaatnnatt ctatttnaat gttgtttcag agtcacagca cagactgaaa ctttttttga 240  
atacctnaat atcacacttn tnccttnaat gatgttgaag acaatgatga catgccttna 300  
gcatataatg tcgac 315

&lt;210&gt; 23

&lt;211&gt; 202

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

actaatccag tgggtgnaa ttccattgtg ttgggcaact caggatatta aatttatnat 60

ttaaaaattc ccaagagaaa naaactccag gccctgattg ttctactggg gaattttacc 120  
aaatgttnca nnaaganatg acgctgattc tgnnaaatct ttttcagaag atagaggaga 180  
acaccaccg nttcatttta tg 202

<210> 24

<211> 365

<212> DNA

<213> Homo sapiens

<400> 24

ggattttcttg cctttttctc cctttttaag tatcaatgta tgaaatccac ctgtaccacc 60  
ctttctgccca tacaaccgct accacatctg gctcctagaa cctgttttgc tttcatagat 120  
ggatctcggga accnagtgtt nacttcattt ttaaacccca ttttagcaga tngtttgctn 180  
tggtctgtct gtattcacca tggggcctgt acacaccacg tgtggttata gtcaaacaca 240  
gtgccctcca ttgtggccac atgggagacc catnaccna tactgcattc tgggctgatn 300  
acggcactgg atctnaccg acntgggatt gaaccgggg tgggcagcng aattgaacag 360  
gatca 365

<210> 25

<211> 359

<212> DNA

<213> Homo sapiens

<400> 25

gtttcctgct tcaacagtgc ttggacggaa cccggcgctc gttccccacc cgggcgggcc 60  
gcccatagcc agccctcgt cactcttca ccgcaccctc ggactgcccc aaggcccccg 120  
ccgcncctcc ngcgccnccg agccaccgcc gcncncncca cctctccttn gtccgcncnt 180  
nacaacgcgt ccaacctgca ngttcgccng aactaccacc nggacteata ngcgcacctc 240  
aaccgcccga tcaacctgga gctctncccc ccgacnttaa cctttccntg tcttacttac 300  
nttaaccgcc gnttattttg cttnaaaaga acttttcccc aatactttct ttcaccnnt 359

<210> 26

<211> 400

<212> DNA

<213> Homo sapiens

<400> 26

agtgaacag tatatgtgaa aaggagttag tgannagcta cataaaaata ttagatatct 60  
ttataatttc caataggata ctcatcagtt ttgaataana gacatattct agagaaacca 120  
ggtttctggt ttcagatttg aactctcaag agcttggaag ttatcactcc catcttcacg 180  
acnacnaana aatctnaacn aacngaana caatgacttt tcttagatct gtcaaagaac 240  
ttcagccacg aggaaaacta tcncctnaa tactggggac tggaaagaga gggtagacag 300  
aatcacagt aatcatagcc caagatcagc ttgcccgag ctnaagctng tacgatnatt 360  
acttacagg accacttcac agtnngtnga tnaantgccn 400

<210> 27

<211> 366

<212> DNA

<213> Homo sapiens

<400> 27

gaatttctta gaaactgaag ttactctgt tccaagatat atcttcactg tcttaatcaa 60  
agggcgctng aatcatagca aatattctca tctttcaact aactttaagt agtntcctg 120  
gaattttaca tttccagaa aacactcctt tctgtatctg tgaaagaaag tgtgcctcag 180  
gctgtagact gggctgcact ggacacctgc gggggactct ggctnagtn ggacatggtc 240

agtattgatt ttctctanac tcagcctgtg tagctntgaa agcatggaac agattacact 300  
gcagttnacg tcatccaca catcttggac tccnagaccc ggggagggtca catagtccgt 360  
tatgna 366

<210> 28

<211> 402

<212> DNA

<213> Homo sapiens

<400> 28

agtgggagcc ttctccttcc cactcagtt ctttacatcc ccgaggcgca gctgggcnaa 60  
ggaagtggcc agctgcagcg cctcctgcag gcagccaacg ttcttgctg tggcctgtgc 120  
agacacatcc ttgccaccac ctttaccgtc catcangcct gacaectgct gcacccactc 180  
gctngctttt aagccccgat nggctgcatt ctgggggact tgacacaggc nctgtatctt 240  
gccagcctca ttgtccaccg tgaagagcat ggcaaaaagt ctgaggggag tgcattctga 300  
anagcttcaa ggcttcattc agggccttng ctnaggcgcc nctctccatc tccnggaata 360  
acnagaggct ggtnnnggtn actntcaata aactgcttcg tc 402

<210> 29

<211> 175

<212> DNA

<213> Homo sapiens

<400> 29

cggacgggca tgaccggtcc ggtcagctgg gtggccagtt tcagttcttc agcagaactg 60  
tctccttct tggggggcga gggcttctct ggggaagagga tgagtttga gcggtactcc 120  
ttcagccgct gcacgttggc ctgcagggac tccgtggact tgttccgct cctcg 175

<210> 30

<211> 360

<212> DNA

<213> Homo sapiens

<400> 30

ttgtatttct tatgatctct gatgggttct tctcgaaaat gccaaagtga agactttgtg 60  
gcagtctcca gatttaaatc cagctgagggc tccctttgtt tttagttcca tgtaacaate 120  
tggaaggaaa cttcacggac aggaagactg ctggagaaga gaagcgtgtt agcccatttg 180  
aggtctgggg aatcatgtaa agggtaacca gacctcactt tttagttattt acatcaatga 240  
gttctttcag ggaaccaaac ccagaattcg gtgcaaaagc caaacatctt ggtgggattt 300  
gataaatgcc ttgggactcg gagtgcctggg cttgtgcaca ggaagagcac cagccgctga 360

<210> 31

<211> 380

<212> DNA

<213> Homo sapiens

<400> 31

acgctctaag cctgtccacg agctcaatag ggaagcctgt gatgactaca gactttgcga 60  
acgctacgcc atggtttatg gatacaatgc tgcctataa cgtacttca ggaagcgccg 120  
agggacnaa tgagctgag ggaagaaaaa aaatctcttt ttttctggag gctggcacct 180  
gattttgtat cccctgttn cagcattncn gaaatacata ggcttatata caatgcttct 240  
ttcctgtata ttctctgtc ttgctgcacc ccttnttccc gccccagat tgataagtaa 300  
tgaaagtga ctgcagtnag ggtcaangga gactcancat atgtgattgt tccntnaaa 360  
acttctggtg tgatactttc 380

<210> 32  
<211> 440  
<212> DNA  
<213> Homo sapiens

<400> 32  
gtgtatggga gccctgact cctcacgtgc ctgatctgtg cccttgggtcc caggtcaggc 60  
ccacccctg cacctccacc tgccccagcc cctgcctctg ccccaagtgg ggccagctgc 120  
cctcacttct ggggtggatg atgtgacctt cctnngggga ctgcggaagg gacaaggggt 180  
ccctgaagtc ttacgggtcca acatcaggac caagtcccat ggacatgctg acaggggtccc 240  
caggggagac cgtntcanta gggatgtgtg cctggctgtg tacgtgggtg tgcagtgcac 300  
gtganaagca cgtggcggtt tctgggggcc atgtttgggg aaggaagtgt gcccnccacc 360  
cttggaagac ctacgtcccn gttagccccct gccctggcac agcngcatnc acttcaaggg 420  
caccctttgg ggggtgggtg 440

<210> 33  
<211> 345  
<212> DNA  
<213> Homo sapiens

<400> 33  
tattttaaca atgtttatta ttcattttatc cctctataga accaccaccc acaccgagga 60  
gattatttgg agtgggtccc aacctagggc ctggactctg aaatctaact cccacttcc 120  
ctcattttgt gacttaggtg ggggcatggt tcagtcagaa ctggtgtctc ctattggatc 180  
gtgcagaagg aggacctagg cacacacata tggtggccac acccaggagg gttgattggc 240  
aggctggaag acaaaagtct cccaataaag gcacttttac ctcaaagang ggggtgggag 300  
tggtctgctg ggaaatgtgt tgttgggggtg ggaagantt atttc 345

<210> 34  
<211> 440  
<212> DNA  
<213> Homo sapiens

<400> 34  
tgtaattttt ttattggaaa acaaatatac aacttgaat ggattttgag gcaaattgtg 60  
ccataagcag attttaagt gctaaacaaa gttaaaaaag caagtaacaa taaaagaaaa 120  
tgtttctggt acaggaccag cagtacaaaa aaatagtgtg cgagtacctg gataatacac 180  
ccgttttgca atagtgaac ttttaagtac atattgttga ctgtccatag tccacgcaga 240  
gttacaactc cacacttcaa caacaacatg ctgacagttc ctaaagaaaa ctactttaaa 300  
aaaggcataa ccagatgtt ccttcatttg accaactcca tctnagtta gatgtgcaga 360  
agggtttana ttttcccaga gtaagccnca tgcaacatgt tacttgatca attttctaaa 420  
ataaggtttt aggacaatga 440

<210> 35  
<211> 540  
<212> DNA  
<213> Homo sapiens

<400> 35  
atagatggaa tttattaagc ttttcacatg tgatagcaca tagttttaat tgcattccaaa 60  
gtactaacaa aaactctagc aatcaagaat ggcagcatgt tattttataa caatcaacac 120  
ctgtggcttt taaaatttgg ttttcataag ataatttata ctgaagtaaa tctagccatg 180  
cttttaaaaa atgcttttagg tctctccaag ctggcagtt aacatttggc ataaacaata 240  
ataaaacaat cacaatttaa taaataacaa atacaacatt gtaggccata atcatatata 300  
gtataaggga aaaggtggta gtgttganta agcagttatt agaatagaat accttggcct 360



ctatgcaa atgtctagac actttgattc actcagccct gacattcagt ttccaaagtt 420  
aggaacagg ttctacagta tcattttaca gtttccaaca cattgaaaac aagtagaaaa 480  
tgatganttg atttttatta atgcattaca tcttcaagan ttatcaccaa cccctcaggt 540

&lt;210&gt; 36

&lt;211&gt; 555

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 36

cttcgtgtgc ttgaaaattg gagcctgccc ctgggcccac aagcccttgc tgggaactga 60  
gaagtgtata tggggcccaa nctactggtg ccagaacaca gagacagcag cccantgcaa 120  
tgctgtcgag cattgcaaac gccatgtgtg gaactaggag gaggaatatt ccactctggc 180  
agaaaccaca gcattggttt ttttctactt gtgtgtctgg gggaatgaac gcacagatct 240  
gtttgacttt gttataaaaa tagggctccc ccacccccc cntttctgtg tnccttattg 300  
tagcantgct gtctgcaagg gagcccttan cccctggcag acanancctgc ttcagtgcgc 360  
ctttctcttc tgctaaatgg atgttgatgc actggaggctc ttttancctg cccttgcagc 420  
gcncctgctg gaggaagana aaactctgct ggcctgaccc acagtttctt gactggangc 480  
cntcaaccct cttgggtgaa gccttgttct gaccctgaca tntgcttggg cncctgggtg 540  
gnctgggctt ctnaa 555

&lt;210&gt; 37

&lt;211&gt; 280

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 37

ccaccgacta taagaactat gccctcgtgt attcctgtac ctgcacatc caactttttc 60  
acgtggattt tgcttggatc ttggcaagaa accctaactc ccctccagaa acagtggact 120  
ctctaaaaaa tatcctgact tctaataaca ttgatntcaa gaaaatgacg gtcacagacc 180  
agggtgaactg ccccnagctc tcgtaaccag gttctacagg gaggtgcac ccactccatg 240  
ttncctctgc ttcgctttcc cctacccccc cccccccat 280

&lt;210&gt; 38

&lt;211&gt; 303

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 38

catcgagctg gttgtcttct tgcttgcctt gtgtcgtaaa atgggggtcc cttactgcat 60  
tatcaaggga aaggcaagac tgggacgtct agtccacagg aagacctgca ccactgtcgc 120  
cttcacacag gtgaactcgg aagacaaagg cgctttggct nagctggtgn aagctatcag 180  
gaccaattac aatgacngat acgatnagat ccgccntcac tggggtagca atgtcctggg 240  
tcctaagtct gtggctcgta tcgccnagct cgaanaggcn aangctaaag aacttgccac 300  
taa 303

&lt;210&gt; 39

&lt;211&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 39

gactcagcgg ctgggtctct tctgtgcac aagcccagca ctccaggctc caaggcattt 60  
atcaaattcc accaagatnt ttggcttttg caccgaattc tgggtttggt tccctnaaag 120  
aactcattga tgtaaatnac tnaaagttag gtctgggtac cctttacatg attccccaga 180

cctcanatgg gctaacacgc ttctcttctc cagcagtctt cctntccgtg aagttacctt 240  
ccagattggt acatggaact gaanacaaag ggagcctcag ctngatttaa atctggagca 300

<210> 40

<211> 318

<212> DNA

<213> Homo sapiens

<400> 40

cccaacacaa tggctgagga caaatcagtt ctctgtgacc agacatgaga aggttgccaa 60  
tgggctggtg ggcgaccaag gccttcccgg agtcttcgtc ctctatgagc tctcgcccat 120  
gatggtgaag ctgacggaga agcacaggtc cttcacccac ttcttgacag gtgtgtgagc 180  
catcattggg ggcattgtca cagtggctgg actcatcgat tcgctcatct accactcagc 240  
acgagccatc cagaaaaaaa ttgatctnng gaagacnacg tagtcaccct cggtncttcc 300  
tctgtctctt ctttctcc 318

<210> 41

<211> 302

<212> DNA

<213> Homo sapiens

<400> 41

acttagatgg ggtccgttca ggggatacca gcgttcacat ttttcctttt aagaaagggg 60  
cttggcctga atgttcccca tccggacaca ggctgcatgt ctctgtnagt gtcaaagctg 120  
ccatnaccat ctcggtaacc tactcttact ccacaatgtc tatnttact gcagggtctt 180  
ataatnagtc cataatgtaa atgcctggcc caagacntat ggctgagtt tatccnaggc 240  
ccaaacnatt accagacatt cctcttanat tgaaaacgga tntctttccc ttggcaaaga 300  
tc 302

<210> 42

<211> 299

<212> DNA

<213> Homo sapiens

<400> 42

cttaataagt ttaaggccaa ggcccgttcc attcttctag caactgacgt tgccagccga 60  
ggtttgagaca tacctcatgt aaatgtggtt gtcaactttg acattcctac ccattccaag 120  
gattacatcc atcgagtagg tcgaacagct agagctgggc gtcgccgaaa ggctattact 180  
tttgtcacac agtatgatgt ggaactcttc cagcgcatag aacacttnat tgggaagaaa 240  
ctaccagggt ttccaacaca ggatgatgag gttatgatgc tnacggaacg cgtcgctna 299

<210> 43

<211> 305

<212> DNA

<213> Homo sapiens

<400> 43

ccaacaatgt caagacagcc gtctgtgaca tcccacctcg tggcctcaan atggcagtca 60  
ccttcattgg caatagcaca gccntccggg agctcttcaa gcgcatctcg gaggagtcca 120  
ctgccatgtt ccgccggaag gccttcctcc actggtacac aggcgagggc atggacaaga 180  
tggagttcac cgaggctgag agcaacatga acgacctcgt ctctnagtat cagcagtacc 240  
gggatgccac cgcagaaana ggaggaggat ttcggtnagg aggccgaaga aggaggcctg 300  
aggca 305

<210> 44

<211> 399  
<212> DNA  
<213> Homo sapiens

<400> 44

```
tttctgtggg ggaaacctga tctcgacnaa attagagaat tttgtcagcg gtatttcggc 60
tggaacagaa cgaaaacnga tnaatctctg tttcctgtat taaagcaact cgatnccag 120
cagacacagc tccnaattga ttccttcttt ngattagcac aacagggaga aagaanatgc 180
ttaacgtatt aagagccnga gactaaacag agctttgaca tgtatgctta ggaaagagaa 240
agaagcagcn gcccgcgnaa ttngaagcng tttctgttgc cntgganaaa gaatttgagc 300
ttctttatta ggccaacgaa aaaccccgaa ananaggcnt tacnatacct tngaaaantc 360
tccngccnna aaaagaaaga agctttcnga ttcttaacc 399
```

<210> 45  
<211> 440  
<212> DNA  
<213> Homo sapiens

<400> 45

```
gcgggagcag aagctaaagc caaagcccaa gagagtggca gtgccagcac tggtgccagt 60
accagtacca ataacagtgc cagtgccagt gccagcacca gtggtggctt cagtgtctgt 120
gccagcctga ccgccactct cacatttggg ctcttcgctg gccttggtgg agctgggtgcc 180
agcaccagtg gcagctctgg tgcttgggt tctctctaca agtgagattt taggtatctg 240
ccttggtttc agtggggaca tctggggcct anggggcnng gataaggagc tggatgatgc 300
taggaaggcc cangttggag aangatgtgn anagtgtgcc aagacactgc ttttggcatt 360
ttattccttt ctgtttgctg gangtcaatt gacccttnna ntttctctta cttgtgtttt 420
canatatngt taatcctgcc 440
```

<210> 46  
<211> 472  
<212> DNA  
<213> Homo sapiens

<400> 46

```
gctctgtaat ttcacatttt aaaccttccc ttgacctcac attcctcttc ggccacctct 60
gtttctctgt tctcttccac agcaaaaact gttcaaaaga gttgttgatt actttcattt 120
ccactttctc acccccatte tcccctcaat taactctcct tcatcccat gatgccatta 180
tgtggtntt attanagtca ccaaccttat tctccaaaac anaagcaaca aggactttga 240
cttctcagca gcactcagct ctggttcttg aaacaccccc gttacttgct attcctecta 300
cctcataaca atctccttcc cagcctctac tgctgccttc tctgagttct tcccagggtc 360
ctaggctcag atgtagtgtg gctcaaccct gctacacaaa gnaatctcct gaaagcctgt 420
aaaaatgtcc atnctgtgcc tgtgagtgtat ctncangna naataacaaa tt 472
```

<210> 47  
<211> 550  
<212> DNA  
<213> Homo sapiens

<400> 47

```
ccttcctccg cctggccatc cccagcatgc tcatgctgtg catggagtgg tgggcctatg 60
aggtcgggag cttctcagc ggtctgtatg aggatggatg acggggactg gtgggaacct 120
gggggccctg tctgggtgca aggcgacagc tgtctttctt caccaggcat cctcggcatg 180
gtggagctgg gcgctcagtc catcgtgtat gaactggcca tcattgtgta catggtccct 240
gcaggcttca gtgtggctgc cagtgtccgg gtangaaacg ctctgggtgc tggagacatg 300
gaagcaggca cggaagtccct ctaccgttct cctgctgatt acagtgtctt ttgctgtanc 360
```

cttcagtgtc ctgctgttaa gctgtaagga tcacntgggg tacattttta ctaccgaccg 420  
agaacatcat taatctgggtg gctcaggtgg ttccaattta tgctgtttcc cacctctttg 480  
aagctcttgc tgctcaggta cagccaatt ttgaaaagta aacaacgtgc ctcggagtgg 540  
gaattctgct 550

&lt;210&gt; 48

&lt;211&gt; 214

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 48

agaaggacat aaacaagctg aacctgcccc agacgtgtga taccagcttc tcagatccag 60  
acaacctcct caacttcaag ctggatcatc gtcctgatna gggcttctac nagagtggga 120  
agtttgtgtt cagttttaag gtgggccagg gttacccgca tgatccccc aaggtgaagt 180  
gtgagacnat ggtctatcac cccnacattg acct 214

&lt;210&gt; 49

&lt;211&gt; 267

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 49

atctgcctaa aatttattca aataatgaaa atnaatctgt ttaagaaat tcagtctttt 60  
agtttttagg acaactatgc acaaatgtac gatggagaat tctttttgga tnaactctag 120  
gtngaggaaac ttaatccaac cggagctntt gtgaaggta gaanacagga gagggaaatc 180  
tggcaaggaa tggagacnga gtttgcaaat tgcagctaga gttaatngtt ntaaatggga 240  
ctgctnttgt gtctccang gaaagtt 267

&lt;210&gt; 50

&lt;211&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 50

gactgggtca aagctgcatg aaaccaggcc ctggcagcaa cctgggaatg gctggaggtg 60  
ggagagaacc tgacttctct tccctctcc ctcctccaac attactggaa ctctgtcctg 120  
ttgggatctt ctgagcttgt tccctgctg ggtgggacag aggacaaagg agaaggaggg 180  
gtctagaaga ggcagccctt ctttgcctc tggggtnaat gagcttgacc tanagtagat 240  
ggagagacca anagectctg atttttaatt tccataanat gttcnaagta tatnttacc 300

&lt;210&gt; 51

&lt;211&gt; 300

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 51

gggtaaaatc ctgcagcacc cactctggaa aatactgctc ttaattttcc tgaaggtggc 60  
ccctatttc tagttgggtc aggattagg atgtggggta tagggcattt aaatcctctc 120  
aagcgtctc caagcacc cggcctgggg gtnagtttct catcccgtta ctgctgctgg 180  
gatcaggttn aataaatgga actcttctg tctggcctcc aaagcagcct aaaaacttag 240  
gggctctgtt agaggggacc tccaccctnn ggaagtcga ggggctnggg aagggtttct 300

&lt;210&gt; 52

&lt;211&gt; 267

&lt;212&gt; DNA

<213> Homo sapiens

<400> 52

aaaatcaact tcntgcatta atanacanat tctanancag gaagtgaana taattttctg 60  
cacctatcaa ggaacnnact tgattgcctc tattnaacan atatategag ttnctatact 120  
tacctgaata ccnccgcata actctcaacc nanatnctc nccatgacac tcnttcttna 180  
atgctantcc cgaattcttc attatatcng tgatgttcgn cctgtnnata tatcagcaag 240  
gtatgtncnn taactgccga nncaang 267

<210> 53

<211> 401

<212> DNA

<213> Homo sapiens

<400> 53

agscttttagc atcatgtaga agcaaaactgc acctatggct gagatagggtg caatgaccta 60  
caagattttg tgttttctag ctgtccagga aaagccatct tcagtcttgc tgacagtcaa 120  
agagcaagtg aaaccatttc cagcctaaac tacataaaaag cagccgaacc aatgattaaa 180  
gacctctaag gctccataat catcattaaa tatgcccāaa ctcatgtga ctttttattt 240  
tatatacagg attaaaatca acattaaatc atcttattta catggccatc ggtgctgaaa 300  
ttgagcattt taaatagtac agtaggctgg tatacattag gaaatggact gcactggagg 360  
caaatagaaa actaaagaaa ttagataggc tggaaatgct t 401

<210> 54

<211> 401

<212> DNA

<213> Homo sapiens

<400> 54

cccaacacaa tggataaaaa cacttatagt aaatggggac attcactata atgatctaag 60  
aagctacaga ttgtcatagt tgttttctg ctttacaaaa ttgtccaga tctggaatgc 120  
cagtttgacc tttgtcttct ataatafttc ctttttttcc cctctttgaa tctctgtata 180  
tttgattctt aactaaaatt gttctcttaa atattctgaa tctctggtat taaaagtgtg 240  
ggtgtatttt ctttacctcc aaggaaagaa ctactagcta caaaaaatat tttggaataa 300  
gcattgtttt ggtataagggt acatattttg gttgaagaca ccagactgaa gtaaacagct 360  
gtgcatccaa tttattatag ttttgtaagt aacaatatgt a 401

<210> 55

<211> 933

<212> DNA

<213> Homo sapiens

<400> 55

tttactgctt ggcaaagtac cctgagcatc agcagagatg ccgagatgaa atcagggaac 60  
tcctagggga tgggtcttct attacctggg aacacctgag ccagatgcct tacaccacga 120  
tgtgcatcaa ggaatgcctc cgcctctacg caccggtagt aaactatccc ggttactcga 180  
caaaccatc acctttccag atggaacgctc cttacctgca ggaataactg tgtttatcaa 240  
tatttgggct cttcaccaca acccctattt ctgggaagac cctcaggtct ttaaccctt 300  
gagattctcc agggaaaatt ctgaaaaaat acatccctat gccttcatac cattctcagc 360  
tggattaagg aactgcattg ggcagcattt tgccataatt gagtgtaaag tggcagtggc 420  
attaactctg ctccgcttca agctggctcc agaccactca aggccacca gctgtcgtca 480  
agttgcctca agtccaagaa tggatccat gtgtttgcaa aaaaagtgtg ctaattttaa 540  
gtccttttctg taaagaatt aakgagacaa ttttcttacc aaaggaagaa caaaaggata 600  
aatataatc aaaaatatg tatatggtg tttgacaaat tatataactt aggatacttc 660  
tgactgggtt tgacatccat taacagtaat ttttaattct ttgctgtatc tggtgaaacc 720

cacaaaaaca cctgaaaaaa ctcaagctga gttccaatgc gaagggaaat gattggtttg 780  
 ggtaactagt ggtagagtgg ctttcaagca tagtttgatc aaaactccac tcagtatctg 840  
 cattactttt atctctgcaa atatctgcat gatagcttta ttctcagtta tctttcccca 900  
 taataaaaaa tatctgcaa aaaaaaaaaa aaa 933

<210> 56

<211> 480

<212> DNA

<213> Homo sapiens

<400> 56

ggctttgaag cttttttgtc tgtgctccct gatcttcagg tcaccaccat gaagttctta 60  
 gcagtcctgg tactcttggg agtttccatc tttctggtct ctgccagaa tccgacaaca 120  
 gctgctccag ctgacacgta tccagctact ggtcctgctg atgatgaagc ccctgatgct 180  
 gaaaccactg ctgctgcaac cactgcgacc actgctgctc ctaccactgc aaccaccgct 240  
 gcttctacca ctgctcgtaa agacattcca gttttaccga aatgggttgg ggatctcccg 300  
 aatggtagag tgtgtccctg agatggaatc agcttgagtc ttctgcaatt ggtcacaact 360  
 attcatgctt cctgtgattt catccaacta cttaccttgc ctacgatac ccttttatct 420  
 ctaatcagtt ttttttcttt caaataaaaa ataactatga gcaacaaaaa aaaaaaaaaa 480

<210> 57

<211> 798

<212> DNA

<213> Homo sapiens

<400> 57

agcctacctg gaaagccaac cagtcctcat aatggacaag atccaccagc tcttctctgtg 60  
 gactaacttt gtgatatggg aagtgaatat agttaacacc ttgcacgacc aaacgaacga 120  
 agatgaccag agtactctta accccttaga actgtttttc cttttgtatc tgcaatatgg 180  
 gatggtattg ttttcatgag cttctagaaa tttcacttgc aagtttattt ttgttctctg 240  
 tgttactgcc attcctattt acagtatatt tgagtgaatg atttatattt taaaaagtta 300  
 catggggctt ttttggttgt cctaaactta caaacattcc actcattctg tttgtaactg 360  
 tgattataat ttttgtgata atttctggcc tgattgaagg aaatttgaga ggtctgcatt 420  
 tatatatatt aaatagattt gataggtttt taaattgctt tttttcataa ggtattttata 480  
 aagttatttg ggggtgtctg ggatttgtgt aaagaaaatt agaaccctgc tgtattttaca 540  
 tttaccttgg tagtttattt gtggatggca gttttctgta gttttgggga ctgtggtagc 600  
 tcttggattg ttttgcaaat tacagctgaa atctgtgtca tggattaaac tggcttatgt 660  
 ggctagaata ggaagagaga aaaaatgaaa tggttgttta ctaattttat actcccatta 720  
 aaaattttta atgttaagaa aaccttaaat aaacatgatt gatcaatatg gaaaaaaaaa 780  
 aaaaaaaaaa aaaaaaaaaa 798

<210> 58

<211> 280

<212> DNA

<213> Homo sapiens

<400> 58

ggggcagctc ctgaccctcc acagccacct ggtcagccac cagctggggc aacgaggggtg 60  
 gaggtccac tgagcctctc gcctgcccc gcactcgtc tgggtgcttg tgatccaagt 120  
 cccctgctg gtccccaca aggactccca tccaggcccc ctctgcccgt ccccttgctca 180  
 tggaccatgg tctgtaggaa gggctcatgc cccttattta tgggaacctt ttcattctaa 240  
 cagaataaac cgagaaggaa accagaaaaa aaaaaaaaaa 280

<210> 59

<211> 382

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 59

```
aggcgggagc agaagctaaa gccaaagccc aagagagtgg cagtgccagc actggtgccca 60
gtaccagtac caataacagt gccagtgccca gtgccagcac cagtgggtggc ttcagtgtctg 120
gtgccagcct gaccgccact ctcacatttg ggctcttcgc tggccttggg ggagctgggtg 180
ccagcaccag tggcagctct ggtgcctgtg gtttctccta caagtgagat tttagatatt 240
gttaatcctg ccagtctttc tcttcaagcc aggggtgcac ctcagaaacc tactcaacac 300
agcactctag gcagccacta tcaatcaatt gaagttgaca ctctgcatta aatctatttg 360
ccattaaaaa aaaaaaaaaa aa 382
```

&lt;210&gt; 60

&lt;211&gt; 602

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 60

```
tgaagagccg cgcggtggag ctgctgcccg atgggactgc caaccttgcc aagctgcagc 60
ttgtggtgga gaatagtgcc cagcgggtca tccacttgcc gggtcagtgg gagaagcacc 120
gggtcccatc ctctgagta ccgccactcc gaaagctgca ggattgcaga gagctggaat 180
cttctcgacg gctggcagag atccaagaac tgcaccagag tgcctgggag gctgctgaag 240
aggcccgcag gaaggaggag gtctataagc agctgatgtc agagctggag actctgcccc 300
gagatgtgtc ccggtgggcc tacaccagc gcctctgga gatcgtgggc aacatccgga 360
agcagaagga agagatcacc aagatcttgt ctgatacga ggagcttcag aaggaaatca 420
actccctatc tgggaagctg gaccggacgt ttgcggtgac tgatgagctt gtgttcaagg 480
atgccaagaa ggacgatgct gttcggaagg cctataagta tctagctgct ctgcacgaga 540
actgcagcca gctcatccag accatcgagg acacaggcac catcatgcgg gaggttcgag 600
ac 602
```

&lt;210&gt; 61

&lt;211&gt; 1368

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 61

```
ccagtgagcg cgcgtaatac gactcaactat agggcgaatt gggtagcggg cccccctcg 60
agcggccgcc cttttttttt tttttttatt gatcagaatt caggctttat tattgagcaa 120
tgaaaacagc taaaacttaa ttccaagcat gtgtagttaa agtttgcaaa gtgggatatt 180
gttcacaaaa cacattcaat gttaaacac tatttatttg aagaacaaaa tatatttaaa 240
attgtttgct tctaaaaagc ccatttccct ccaagtctaa actttgtaat ttgatattaa 300
gcaatgaagt tattttgtac aatctagtta aacaagcaga atagcactag gcagaataaa 360
aaattgcaca gacgtatgca attttccaag atagcattct ttaaattcag ttttcagctt 420
ccaaagattg gttgcccata atagacttaa acatataatg atggctaaaa aaaataagta 480
tacgaaaatg taaaaaagga aatgtaagtc cactctcaat ctcataaaag gtgagagtaa 540
ggatgctaada gcaaaataaa tgtaggttct tttttctgt ttcggtttat catgcaatct 600
gcttctttga tatgccttag ggttacccat ttaagttaga ggttgtaatg caatggtggg 660
aatgaaaatt gatcaaatat acaccttgtc atttcatttc aaattgcggg ctggaaactt 720
ccaaaaaag ggtaggcatg aagaaaaaaa aaatcmaatc agaactctct caggggtttg 780
kgktctgata tggcagacar gatacaagtc ccaccaggag atggagcaat tcaaaaataag 840
ggtaatgggc tgacaaggta ttattgccag catgggacag aatgagcaac aggctgaaaa 900
gtttttggat tatatagcac ctagagtctc tgatgtaggg aatttttggt agtcaaacat 960
acgctaaact tccaaggga aatctttcag gtacgctaag ctgtcttttc tagagtgatg 1020
agttgcattg ctactgtgat tttttgaaaa caaactgggt ttgtacaagt gagaaagact 1080
agagagaaaag attttagtct gtttagcaga agccatttta tctgcgtgca catggatcaa 1140
```

```
tatttctgat cccctatacc ccaggaaggg caaaatccca aagaaatgtg ttagcaaaat 1200
tggctgatgc tatcatattg ctatggacat tgatcttgcc caacacaatg gaattccacc 1260
acactggact agtggatcca ctagtcttag agcggccggc caccgcggtg gagctccagc 1320
ttttgttccc tttagtggag gttaattgcg cgcttggcgt aatcatnn 1368
```

&lt;210&gt; 62

&lt;211&gt; 924

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 62

```
caaaggnaca ggaacagctt gnaaagtact gncatnccn cctgcagga ccagcccttt 60
gcctccaaaa gcaataggaa atttaaaaga tttnactga gaaggggncc acgttttntart 120
tntnaatgtn tcargnanar tnccttncaa atgncrntn cactnactnr gnatttggtt 180
tnccgnrtnc mgnactatnt caggtttgaa aaactggatc tgccacttat cagttatgtg 240
accttaaaga actccgttaa tttctcagag cctcagtttc cttgtctata agttgggagt 300
aatattaata ctatcatttt tccaaggatt gatgtgaaca ttaatgaggt gaaatgacag 360
atgtgtatca tggttcctaa taacatcca aaatatagta cttactattg tcattattat 420
tacttgtttg aagctaaaga cctcacaata gaatcccatc cagcccacca gacagagytc 480
tgagttttct agtttggaag agctattaaa taacaacktc tagtgtcaat tctatacttg 540
ttatggtcaa gtaactgggc tcagcatttt acattcattg tctctttaag ttctagcaat 600
gtgaagcagg aactatgatt atattgacta cataaatgaa gaaattgagg ctacagataca 660
ttaagtaatt ctcccagggt cacacagcta gaactggcaa agcctgggat tgatccatga 720
tcttccagca ttgaagaatc ataaatgtaa ataaactgaa ggccttttcc tcagaagagc 780
tcctgggtgct tgcaccaacc cactagcact tgttctctac aggggaacat ctgtgggcct 840
gggaatcact gcacgtcgca agagatgttg cttctgatga attattgttc ctgtcagtgg 900
tgtgaaggca aaaaaaaaaa aaaa 924
```

&lt;210&gt; 63

&lt;211&gt; 1079

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 63

```
agtcccaaga actcaataat ctcttatgtt ttcttttgaa gacttatttt aaatattaac 60
tatttcggtg cctgaatgga aaaatataaa cattagctca gagacaatgg ggtacctgtt 120
tggaatccag ctggcagcta taagcacctg tgaaaactct gacaggcttt gtgccctttt 180
tattaaatgg cctcacatcc tgaatgcagg aatgtgttcg tttaataaaa cattaatctt 240
taatgttgaa ttctgaaaac acaaccataa atcatagtgt gtttttctgt gacaatgatc 300
tagtacatta ttctctccac agcaaaccta ctttccaga aggtggaaat tgtatttgca 360
acaatcaggg caaaaccac acttgaaaag cattttacaa tattatctct aagttgcaca 420
gaagacccca gtgatcacta ggaaatctac cacagtccag tttttctaata ccaagaaggt 480
caaacttcgg ggaataatgt gtccctcttc tgctgctgct ctgaaaaata ttcatcaaa 540
acgaagttaa caagcagcag ttattccaag attagagttc atttgtgtat cccatgtata 600
ctggcaatgt ttagggttgc ccaaaagctc ccagacatcc acaatgttgt tgggtaaacc 660
accacatctg gtaacctctc gatcccttag atttgtatct cctgcaataa taactgtagc 720
tgactctgga gcctcttgca tttcttttaa aaccattttt aactgattca ttcgttcgcg 780
agcatgccct ctggtgctct ccaaatggga tgcataagg caaagctcat ttcctgacac 840
attcacatgc acacataaaa ggttctctat catttttgta cttggaaaag gaataatctc 900
ttggcttttt aatttcaact ttgatttctt caacattata gctgtgaaat atccttcttc 960
atgacctgta ataattctat aattacttga tctcttctt aggtagctat aatatggggg 1020
aataacttcc tgtagaataa tcacatctgg gctgtacaaa gctaagtagg aacacacc 1079
```

&lt;210&gt; 64

&lt;211&gt; 1001



<212> DNA  
 <213> Homo sapiens

<400> 64  
 gaatgtgcaa cgatcaagtc aggggtatctg tggatccac cactttgagc atttatcgat 60  
 tctatatgtc aggaacattt caagttatct gttctagcaa ggaaatataa aatacttata 120  
 gtttaactatg gcctatctac agtgcaacta aaaactagat tttattcctt tccacctgtg 180  
 gggttgattt catttaccac cctcttttca ttccctttct caccacaca ctgtgccggg 240  
 cctcaggcat atactattct actgtctgtc tctgtaagga ttatcatttt agcttccaca 300  
 tatgagagaa tgcattgcaa gtttttcttt ccattgtctg cttatttcac ttaacataat 360  
 gacctccgct tccatccatg ttatttatat tacccaatag tgttcataaa tatatatata 420  
 cacatatata ccacattgca ttgttccaat tattcattga cggaaactgg ttaagtattt 480  
 atcggtgcta ttgtggatag tgcgtcaata aacacgcaag tggggatata attgaagag 540  
 ttttttgggt gatgttctc caaattttta gattgttttg tctatgtttg tgaaaatggc 600  
 gtttagtattt tcatagagat tgcattgaat ctgtagattg ctttgggtta gtatgggtat 660  
 tttgatggta ttaatttttt cattccatga agatgagatg tctttccatt gtttgtgtcc 720  
 tctacatttt ctttcatcaa agttttgttg tatttttgaa gtagatgtat ttcaccttat 780  
 agatcaagtg tattccctaa atattttatt tttgtagcta ttgtatga aattgccttc 840  
 ttgatttctt tttcacttaa ttcattatta gtgtatggaa atgttatgga tttttatttg 900  
 ttggttttta atcaaaaact gtattaaact tagagttttt tgtggagttt ttaagttttt 960  
 ctgatataaa gatcatgaca tctaccaaaa aaaaaaaaaa a 1001

<210> 65  
 <211> 575  
 <212> DNA  
 <213> Homo sapiens

<400> 65  
 acttgatata aaaaggatat ccataatgaa tattttatag tgcattcctt acattagcca 60  
 cttaaatacgt tattgcttga tgaagacctt tcacagaatc ctatggattg cagcatttca 120  
 cttggctact tcataccat gccctaaaga ggggcagttt ctcaaaagca gaaacatgcc 180  
 gccagttctc aagttttcct cctaactcca tttgaatgta agggcagctg gcccccattg 240  
 tggggagggt cgaacatttt ctgaattccc attttcttgt tcgctggctaa atgacagttt 300  
 ctgtcattac ttagattccc gatctttccc aaagggtgtg atttacaag aggccagcta 360  
 atagccagaa atcatgacct tgaaagagag atgaatttc aagctgtgag ccaggcagga 420  
 gctccagtat ggcaagggtt cttgagaatc agccatttgg tacaataaag atttttaag 480  
 cttttatgtt ataccatgga gccatagaaa ggctatggat tgtttaagaa ctattttaaa 540  
 gtgttccaga cccaaaaagg aaaaaaaaaa aaaa 575

<210> 66  
 <211> 831  
 <212> DNA  
 <213> Homo sapiens

<400> 66  
 attgggctcc ttctgctaaa cagccacatt gaaatggttt aaaagcaagt cagatcaggt 60  
 gatttgtaaa attgtattta tctgtacatg tatgggcttt taattccac caagaaagag 120  
 agaaattatc tttttagtta aaaccaaatt tcaactttca aaatatcttc caacttattt 180  
 attggtgtc actcaattgc ctatatatat atatatatat gtgtgtgtgt gtgtgtgcgc 240  
 gtgagcgac gtgtgtgtat gcgtgcgcac gtgtgtgtat gtgtattatc agacataggt 300  
 ttctaacttt tagatagaag aggagcaaca tctatgccaa atactgtgca ttctacaatg 360  
 gtgctaattc cagacctaaa tgatactcca tttaatttaa aaaagagttt taaataatta 420  
 tctatgtgcc tgtatttccc ttttgagtgc tgcacaacat gttacatat tagtgtaaaa 480  
 gcagatgaaa caaccacgtg ttctaaagtc tagggattgt gctataatcc ctatttagtt 540  
 caaaattaac cagaattctt ccattgtgaaa tggaccaaac tcatattatt gttatgtaaa 600

tacagagttt taatgcagta tgacatccca caggggaaaa gaatgtctgt agtgggtgac 660  
 tgttatcaaa tttttatag aatacaatga acggtgaaca gactggtaac ttgtttgagt 720  
 tcccatgaca gatttgagac ttgtcaatag caaatcattt ttgtatttaa atttttgtac 780  
 tgatttgaaa aacatcatta aatatcttta aaagtaaaaa aaaaaaaaaa a 831

<210> 67

<211> 590

<212> DNA

<213> Homo sapiens

<400> 67

gtgctctgtg tttttttta ctgcattaga cattgaatag taatttgcgt taagatacgc 60  
 ttaaaggctc ttgtgacca tgttccctt ttagcaata aaatgtttt tacgaaaact 120  
 ttctccctgg attagcagtt taaatgaac agagttcatc aatgaaatga gtatttaaaa 180  
 taaaaatttg ccttaatgta tcagttcagc tcacaagtat tttaaatga ttgagaagac 240  
 ttgaattaaa gaaaaaaaaa ttctaatca tttttttaa atataagact aaaattgttt 300  
 ttaaacaca ttcaaatag aagtgaagtt gaactgacct tttttatact ctttttaagt 360  
 ttgttccttt tccctgtgcc tgtgtcaat ctcaagtc tgcgtgaaat acatttgata 420  
 caaagtttct ttagttgtg ttagttcttt tgcctgtct gtttttggct gaagaaccaa 480  
 gaagcagact tttcttttaa aagaattatt tctctttcaa atatttctat ccttttttaa 540  
 aaattccttt ttatggctta tatactaca tttttaaaaa aaaaaaaaaa 590

<210> 68

<211> 301

<212> DNA

<213> Homo sapiens

<400> 68

ttgtgttggg gttccctttt ccggtcggcg tggctcttgcg agtggagtgt ccgctgtgcc 60  
 cgggcctgca ccatgagcgt ccggccttc atcgacatca gtgaagaaga tcaggctgct 120  
 gagcttcgtg cttatctgaa atctaaggga gctgagattt cagaagagaa ctcggaaggt 180  
 ggacttcagc ttgatttagc tcaaattatt gaagcctgtg atgtgtgtct gaaggaggat 240  
 gataaagatg ttgaaagtgt gatgaacagt ggggnatcct actcttgatc cggaanccna 300  
 c

<210> 69

<211> 301

<212> DNA

<213> Homo sapiens

<400> 69

tctatgagca tgccaaggct ctgtgggagg atgaaggagt gcgtgcctgc tacgaacgct 60  
 ccaacgagta ccagctgatt gactgtgcc agtacttctt ggacaagatc gacgtgatca 120  
 agcaggctga ctatgtgcc agcgatcagg acctgcttcg ctgccgtgtc ctgacttctg 180  
 gaattcttga gaccaagttc caggtggacn aagtcaactt ccacatgntt gacgtgggtg 240  
 gccagcgcga tgaacgccgc aagtggatcc agtgcttcaa cgatgtgact gccatcatct 300  
 t

<210> 70

<211> 201

<212> DNA

<213> Homo sapiens

<400> 70

gcggctcttc ctccggcagc ggaagcggcg cggcggctcg agaagtggcc taaaacttcg 60

gcgttgggtg aaagaaaatg gcccgaacca agcagactgc tcgtaagtcc accggtggga 120  
 aagccccccg caaacagctg gccacgaaag ccgccaggaa aagcgctccc tctaccggcg 180  
 gggatgaaga gcctcatcgc t 201

&lt;210&gt; 71

&lt;211&gt; 301

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

gccggggtag tcgccgncgc cggcgccgct gcagccactg caggcaccgc tgccgcccgc 60  
 tgagtagtgg gcttaggaag gaagaggtca tctcgctcgg agcttcgctc ggaaggggtct 120  
 ttgttccttg cagccctccc acgggaatga caatggataa aagtgaactg gtacanaaag 180  
 ccaaactcgc tgagcaggct gagcgatatg atgatatggc tgcagccatg aaggcagtcg 240  
 cagaacaggg gcatgaactc ttcaacgaag agagaaatct gctctctggg gcctacaaga 300  
 a 301

&lt;210&gt; 72

&lt;211&gt; 251

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 72

cttggggggg gttgggggag agactgtggg cctggaaata aaacttgtct cctctaccac 60  
 caccctgtac cctagcctgc acctgtccac atctctgcaa agttcagctt ccttccccag 120  
 gtctctgtgc actctgtctt ggatgtcttg gggagctcat ggggtggagg gtctccacca 180  
 gagggaggct caggggactg gttggggccg ggatgaatat ttgagggata aaaatttgtt 240  
 aagagccaan g 251

&lt;210&gt; 73

&lt;211&gt; 913

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 73

tttttttttt tttttccag gccctctttt tatttacagt gataccaaac catccacttg 60  
 caaattcttt ggtctcccat cagctggaat taagtaggta ctgtgtatct ttgagatcat 120  
 gtatttgtct ccactttggt ggatacaaga aaggaaggca cgaacagctg aaaaagaagg 180  
 gtatcacacc gctccagctg gaatccagca ggaacctctg agcatgccac agctgaacac 240  
 ttaaaaggag aaagaaggac agctgtctct catttatttt gaaagcaaat tcatttgaaa 300  
 gtgcataaat ggtcatcata agtcaaacgt atcaattaga ccttcaacct aggaacaaa 360  
 attttttttt tctatttaat aatacaccac actgaaatta ttgccaatg aatcccaag 420  
 atttggtaga aatagtacaa ttctgtattt ctttctctct tcttttcttc agacaaacac 480  
 caaataaaat gcagggtgaaa gagatgaacc acgactagag gctgacttag aaatttatgc 540  
 tgactcgatc taataaaaaat tatgttggtt aatgttaatc tatctaaaat agagcatatt 600  
 gggaatgctt ttcaaagaag gtcaagtaac agtcatacag ctagaaaagt ccctgaaaaa 660  
 aagaattgtt aagaagtata ataacctttt caaaaccac aatgcagctt agttttcctt 720  
 tatttatttg tggcatgaa gactatcccc atttctccat aaaatcctcc ctccatactg 780  
 ctgcattatg gcacaaaaga ctctaagtgc caccagacag aaggaccaga gtttctgatt 840  
 ataaacaatg atgtcgggta atgtttaaat gagaacattg gatatggatg gtcagcccaa 900  
 cacaatggaa ttc 913

<210> 74  
 <211> 351  
 <212> DNA  
 <213> Homo sapiens

<400> 74  
 tgtgcnacagg ggatgggtgg gcngtggaga ngatgacaga aaggctggaa ggaanggggg 60  
 tgggttttgaa ggccanggcc aaggggncct cagggtccgnt tctgnnaagg gacagccttg 120  
 aggaaggagn catggcaagc catagctagg ccaccaatca gattaagaaa nnctgagaaa 180  
 nctagctgac catcactgtt ggtgnccagt ttcccaacac aatggaatnc caccacactg 240  
 gactagnnga nccactagtt ctagagcggc cgccaccgcg gtggaacccc aacttttgcc 300  
 cctttagnga gggtaattg cgcgcttggc ntaatcatgg tcataagctg t 351

<210> 75  
 <211> 251  
 <212> DNA  
 <213> Homo sapiens

<400> 75  
 tacttgacct tctttgaaaa gcattcccaa aatgctctat tttagataga ttaacattaa 60  
 ccaacataat tttttttaga tcgagtcagc ataaatttct aagtcagcct ctatgctgtg 120  
 ttcattctctt tcacctgcat tttatttggg gtttgtctga agaaaggaaa gaggaagca 180  
 aatacgaatt gtactatttg taccaaactt ttgggattca ttggcaaata atttcagtgt 240  
 ggtgtattat t 251

<210> 76  
 <211> 251  
 <212> DNA  
 <213> Homo sapiens

<400> 76  
 tatttaataa tacaccacac tgaaattatt tgccaatgaa tcccaaagat ttggtacaaa 60  
 tagtacaatt cgtatttgct ttctctttc ctttcttcag acaaacacca aataaaatgc 120  
 aggtgaaaga gatgaaccac gactagaggc tgacttagaa atttatgctg actcgatcta 180  
 aaaaaatta tgttgggttaa tgtaaatcta tctaaaatag agcattttgg gaatgctttt 240  
 caaagaaggt c

<210> 77  
 <211> 351  
 <212> DNA  
 <213> Homo sapiens

<400> 77  
 actcacctg ctgtgtgctg tgtgcctgct gcctggcagc ctggccctgc cgctgctcag 60  
 gaggcgggag gcatgagtga gctacagtgg gaacaggctc aggactatct caagagann 120  
 tatctctatg actcagaaac aaaaaatgcc aacagttag agccaaact caaggagatg 180  
 caaaaattct ttggcctacc tataactgga atgttaaact cccgcgtcat agaaataatg 240  
 cagaagccca gatgtggagt gccagatggt gcagaatact cactatttcc aaatagccca 300  
 aaatggactt ccaaagtggc cacctacagg atcgtatcat atactcgaga c 351

<210> 78  
 <211> 1592  
 <212> DNA  
 <213> Homo sapiens

&lt;400&gt; 78

gaattccatt	gtgttggggc	cctggggggc	gaggggaggg	gccaccacg	gccttatttc	60
cgcgagcgc	ggcactgcc	gctccgagcc	cgtgtctgtc	gggtgccgag	ccaactttcc	120
tgcgtccatg	cagccccgc	ggcaacggct	gcccgcctcc	tggtcgggc	ccagggggcc	180
gcgccccacc	gccccgtgc	tgcgctgtc	gctgttgctc	gccccggtg	cgcgccccgc	240
ggggtccggg	gaccccgacg	accctgggca	gcctcaggat	gctgggggtcc	cgcgaggct	300
cctgcagcag	gcggcgcg	cggcgttca	cttcttcaac	ttccggtccg	gctcgccag	360
cgcgtcgcga	gtgctggccg	aggtgcagg	gggcccgcg	tggattaatc	caaaagagg	420
atgtaaagt	cacgtgtct	tcagcacaga	gcgtacaac	ccagagtctt	tacttcagga	480
aggtgaggg	cgtttgggga	aatgttctgc	tcgagtgtt	ttcaagaatc	agaaaccag	540
accaactatc	aatgtaact	gtacacggct	catcgagaaa	aagaaaagac	aacaagagga	600
ttacctgctt	tacaagcaaa	tgaagcaact	gaaaaacccc	ttggaaatag	tcagcatacc	660
tgataatcat	ggacatattg	atccctctct	gagactcatc	tgggatttgg	ctttccttgg	720
aagctcttac	gtgatgtggg	aaatgacaac	acagggtgtc	cactactact	tggcacagct	780
caactagtgtg	aggcagtgg	aaactaatga	tgatacaatt	gattttgatt	atactgttct	840
acttcattgaa	ttatcaacac	aggaaataat	tcctgtgcg	attcacttgg	tctggtacc	900
tggcaaacct	cttaaagtga	agtaaccactg	tcaagagcta	cagacaccag	aagaagcctc	960
cggaaactgaa	gaaggatcag	ctgtagtacc	aacagagctt	agtaatttct	aaaaagaaaa	1020
aatgatcttt	ttccgacttc	taaacaagt	actatactag	cataaatcat	tcttctagta	1080
aaacagctaa	ggtatagaca	ttctaataat	ttgggaaaac	ctatgattac	aagtaaaaac	1140
tcagaaatgc	aaagatgttg	gtttttgtt	tctcagtctg	ctttagcttt	taactctgga	1200
agcgcattgca	cactgaactc	tgtcagtgc	taaacagtca	ccagcagggt	cctcagggtt	1260
tcagccctaa	aatgtaaaac	ctggataatc	agtgtatgtt	gcaccagaat	cagcattttt	1320
tttttaactg	caaaaaatga	tggctctcatc	tctgaattta	tatttctcat	tcttttgaac	1380
atactatagc	taatatattt	tatgttgcta	aattgcttct	atctagcatg	ttaaacaag	1440
ataatatact	ttcgatgaaa	gtaaattata	ggaaaaaat	taactgtttt	aaaaagaact	1500
tgattatgtt	ttatgatttc	aggcaagtat	tcatttttaa	cttgctacct	acttttaaat	1560
aaatgtttac	atttctaaaa	aaaaaaaaaa	aa			1592

&lt;210&gt; 79

&lt;211&gt; 401

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 79

catactgtga	attgttcttg	actccttttc	ttgacattca	gttttcanaa	tttccatctt	60
tcttctggaa	ctaattgtgt	gttctcttga	ctgcctgctg	ggccagcatc	cgattgccag	120
ccagaaacgt	cacactgccc	aagatggcca	ggtacttcaa	ggtctggaac	atgttgagct	180
gagtccagta	gacatacatg	agtcccagca	tagcagcatg	tcccagggtga	aatataatcg	240
tgctaggagc	aaaagtgaag	ttggagacat	tggcaccat	ccggatccac	tagttctaga	300
gcggccgcca	ccgcggtgga	gctccagctt	ttgttccctt	tagtgagggt	taattgcgcg	360
cttgccgtaa	tcatggnat	agctgtttcc	tgtgtgaaat	t		401

&lt;210&gt; 80

&lt;211&gt; 301

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 80

aaaaatgaaa	catctatctt	agcagcaaga	ggctgtgagg	gatggggtag	aaaaggcatc	60
ctgagagagt	tctagaccga	cccaggctct	gtggcacact	atacgggtca	ggaggggtgg	120
aagacaggcc	taagctctag	gacggtgaat	ctcggggcta	tttgtggatt	tgtagaaaac	180
agacattctt	ttggcctttt	cctggcactg	gtgttgccgg	cagggtggga	gaagtgaagg	240
accagtcact	gttcagtcac	tgccaccaca	gatcttcagc	agaatcttcc	ggtaatcccc	300
t						301

<210> 81  
 <211> 301  
 <212> DNA  
 <213> Homo sapiens

<400> 81  
 tagccagggtt gctcaagcta attttattct ttcccaacag gatccatttg gaaaatatca 60  
 agccttttaga atgtggcagc aagagaaagc ggactacgca ggaacgggga gtttgggaga 120  
 agctctcctg gtgttgactt agggatgaag gctccaggct gctgccagaa atggagtcac 180  
 cagcagaaga actgntttct ctgataagga tgtcccaacca ttttcaagct gttcggttaa 240  
 gttacacagg tccttcttgc agcagtaagt accgtagct cattttccct caagcgggtt 300  
 t 301

<210> 82  
 <211> 201  
 <212> DNA  
 <213> Homo sapiens

<400> 82  
 tcaacagaca aaaaaagttt attgaatata aaactcaaag gcatcaacag tcctggggccc 60  
 aagagatcca tggcaggaag tcaagagttc tgcttcaggg tcggtctggg cagccctgga 120  
 agaagtcatt gcacatgaca gtgatgagtg ccaggaaaac agcatactcc tggaaagtcc 180  
 acctgctggn cactgnttca t 201

<210> 83  
 <211> 251  
 <212> DNA  
 <213> Homo sapiens

<400> 83  
 gtaaggagca tactgtgccc atttattata gaatgcagtt aaaaaaata ttttgagggt 60  
 agcctctcca gtttaaaagc acttaacaag aaacacttgg acagcgatgc aatggtctct 120  
 cccaaaccgg ctccctctta ccaagtaccg taaacagggt ttgagaacgt tcaatcaatt 180  
 tcttgatatg aacaatcaaa gcatttaatt caaacatatt tgcttctcaa anaataaaac 240  
 cattttccaa a 251

<210> 84  
 <211> 301  
 <212> DNA  
 <213> Homo sapiens

<400> 84  
 agtttataat gttttactat gatttagggc ttttttttca aagaacaaaa attataagca 60  
 taaaaactca ggtatcagaa agactcaaaa ggctgtttt cactttgttc agattttgtt 120  
 tccaggcatt aagtgtgtca tacagttgtt gccactgctg ttttccaaat gtccgatgtg 180  
 tgctatgact gacaactact tttctctggg tctgatcaat tttgcagtan accattttag 240  
 ttcttacggc gtcnataaca aatgcttcaa catcatcagc tccaatctga agtcttgctg 300  
 c 301

<210> 85  
 <211> 201  
 <212> DNA  
 <213> Homo sapiens

&lt;400&gt; 85

tattttgtgta tgtaacattt attgacatct acccactgca agtatagatg aataagacac 60  
agtcacacca taaaggagtt tatccttaaa aggagtgaag gacattcaaa aaccaactgc 120  
aataaaaaag ggtgacataa ttgctaaatg gagtggagga acagtgttta tcaattcttg 180  
attgggccac aatgatatac c 201

&lt;210&gt; 86

&lt;211&gt; 301

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 86

tttataaaat attttattta cagtagagct ttacaaaaat agtcttaaat taatacaaat 60  
cccttttgcā atātaāctta tatgactatc ttctcaaaaa cgtgacattc gattataaca 120  
cataaactac atttatagtt gttāagtcac cttgtāgtat aaatātgttt tcatcttttt 180  
tttgtāataa ggtacatācc aataacaatg aacaatggac aacaatctt attttgntat 240  
tcttccaatg taaattcat ctctggccaa aacaaaatta accaaagaaa agtaaaacaa 300  
t 301

&lt;210&gt; 87

&lt;211&gt; 351

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 87

aaaaaagatt taagatcata aataggtcat tgttgcaca acacatttca gaatcttaaa 60  
aaaacaaaca ttttggcttt ctaāgāaaaa gacttttaaa aaaaatcaat tccctcatca 120  
ctgaaaggac ttgtacattt ttaāacttcc agtctcctaa ggcacagtat ttaātcagaa 180  
tgccaatatt accaccctgc tgtāgcāga atāāagaagc aagggtāta cacttāaaaa 240  
aacngccaaa ttctgaacc aaatcattgg cattttāaaa aagggtāta aaaaācngnt 300  
āaggggggga gcattttaag taaāgaāngg ccaagggtgg tatgācngga c 351

&lt;210&gt; 88

&lt;211&gt; 301

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 88

gttttaggtc ttaccaatt tgattggtt atcaacaggg catgaggtt aatatatct 60  
ttgaggaaag gtaaagtcaa atttgacttc ataggatcāc ggcgtcctca ctctgtgca 120  
ttttctggtg gaagcacaca gttāattāac tcaagtgtgg cgtāgcgāc gctttttcat 180  
ggngtcattt atccacttgg tgaācttgcā cacttgaatg naaactcctg ggtcattggg 240  
ntggccgcaa gggāaggtc cccaagacac caaaccttgc aggtacctn tgcacacaa 300  
c 301

&lt;210&gt; 89

&lt;211&gt; 591

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 89

tttttttttt tttttttatt aatcaaatga ttcaaaacaa ccatcattct gtcaatgcc 60  
aagcaccag ctggtcctct cccacatgt cacactctcc tcagcctctc ccccaaccct 120  
gctctcctc ctccctgcc ctāgcccāg gacagagtct agggāggāc tggggcāgāg 180  
ctggāggcāg gaagāgāgā ctggācāgāc agctatggtt tggāttggg āgāgāttāg 240

gaagtaggtt	cttaaagacc	cttttttagt	accagatatt	cagccatatt	cccagctcca	300
ttattcaa	at	tttcccat	agcccagctc	ctctctgttc	tccccctact	360
tggtcttac	acaattttta	tccctcaa	at	catccct	ggcccaacca	420
cctccctctg	gtggagactc	ctccacccat	gagctcccca	gagcatccaa	gacagagtgc	480
acagagacct	gggaaggaa	gctgaacttt	gcagagatgt	ggacaggtgc	aggctagggt	540
acaggggtggt	ggtagaggag	acaagtttta	tttccaggcc	cacagtctct	c	591

&lt;210&gt; 90

&lt;211&gt; 1996

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 90

tttttttttt	ttttttatca	aatgaatact	ttattagaga	cataacacgt	ataaaataaa	60
tttctttttca	tcatggagtt	accagatttt	aaaaccaacc	aacactttct	cattttttaca	120
gctaagacat	gttaaattct	taaatgccat	aatttttgtt	caactgcttt	gtcattcaac	180
tcacaagtct	agaatgtgat	taagctacaa	atctaagtat	tcacagatgt	gtcttaggct	240
tggtttgtaa	caatctagaa	gcaatctggt	tacaaaagtg	ccaccaaaagc	attttaaaga	300
aaccaattta	atgccaccaa	acataagcct	gctataacctg	ggaaacaaaa	aatctcacac	360
ctaaattcta	gcagagtaaa	cgattccaac	tagaatgtac	tgtatatcca	tatggcacat	420
ttatgacttt	gtaatatgta	attcataata	cagggttagg	tgtgtggtat	ggagctagga	480
aaaccaaaagt	agtaggatat	tatagaaaag	atctgatgtt	aagtataaag	tcatatgcct	540
gatttcctca	aaccttttgt	ttttcctcat	gtcttctgtc	tttatatttt	tatcacaaac	600
caagatctaa	cagggttctt	tctagaggat	tattagataa	gtaacacttg	atcattaagc	660
acggatcatg	ccactcattc	atggttggtc	tatgttccat	gaactcta	agcccaactt	720
atacatggca	ctccaagggg	atgcttcagc	cagaaagtaa	agggctgaaa	aagtagaaca	780
atacaaaagc	cctcgtgtgg	tgggaactgt	ggcctcactc	ttacttgtcc	ttccattcaa	840
aacagtttgg	cacctttcca	tgacgaggat	ctctacaggt	aggttaaaa	acttttctgt	900
gctattcagc	cagaaatagt	ttttgtgctg	gatatgattt	taaaacagat	tttgtctgtc	960
accagtgc	aaacattaca	gatgtctggg	ctaatacaaa	aacacataag	aatctacaac	1020
tttataattta	atactctatt	caaatttaac	tcaaaagta	gcaaaataat	tagaagttaa	1080
aacttaattc	ttctgagagc	tctatttgga	aaagcttcac	atatccacac	acaaatatgg	1140
gtatatctat	gcacagggca	aacaactgta	ttctgaagca	taaataaact	caaagtaaga	1200
catcagttagc	tagataccag	ttccagattt	ggttaatggg	ctctggggat	cccattttaa	1260
gcactctcag	atgaggatct	tgctcagttg	ttagactatc	attagtttga	ttaagcaact	1320
gaagtttact	tcataaatta	ctttttccta	tatccaggac	tctgcctgag	aaattttata	1380
cattctctca	aaggtaagta	ttctccaaag	gtaagtattt	gactattaac	acaaaggcaa	1440
tgtgattatt	gcataatgac	actaaatatt	atgtggcttt	tctgttaggt	ttataagttt	1500
tcaatgatca	gttcaagaaa	atgcagatca	tatataacta	aggttttaca	ccagtggttg	1560
acaaactatg	gcccacaggc	taaacccagc	ctccccctgt	ttttataaat	aagttttatt	1620
agacataacc	acactcattc	atctctgtat	tgtgtatagc	tgcttttcacg	ctatactagc	1680
agaactgaat	agttgtgaca	gagactgtat	ggaccgtgaa	gcataaata	ttaccatctg	1740
gcccattcta	aaaaaagtgt	gccaattcct	ggtttacact	aaaatataga	gtttagtggg	1800
aagcctattt	gaaatgtgtt	tttttttaggg	gctgtaatta	ccaattaaaa	ttaaggttca	1860
gggtgactcag	caaccaaaaca	aaagggatac	taatttttta	tgaacaatat	atttgtattt	1920
tatggacata	aaaggaaact	ttcagaaaga	aaaggaggaa	aataaagggg	gaaagggacc	1980
caacacaatg	gaattc					1996

&lt;210&gt; 91

&lt;211&gt; 911

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 91

gccctttttt	tttttttttt	cttgttttaa	aaaattgttt	tcattttaat	gatctgagtt	60
------------	------------	------------	------------	------------	------------	----



agtaacaaac	aaatgtacaa	aattgtcttt	cacatttcca	tacattgtgt	tatggaccaa	120
atgaaaacgc	tggactacaa	atgcaggttt	ctttatatcc	ttaacttcaa	ttattgtcac	180
ttataaataa	aggtgatttg	ctaacacatg	catttgtgaa	cacagatgcc	aaaaattata	240
catgtaagtt	aatgcacaac	caagagtata	cactgttcat	ttgtgcagtt	atgcgtcaaa	300
tgcgactgac	acagaagcag	ttatcctggg	atatttcact	ctatatgaaa	agcatcttgg	360
agaaatagat	tgaataacag	tttaaaacaa	aaattgtatt	ctacaaatac	aataaaattt	420
gcaacttgca	catctgaagc	aacatttgag	aaagctgctt	caataaccct	gctgttatat	480
tggttttata	ggtatatctc	caaagtcag	ggttgggata	tagctgcttt	aaagaaaata	540
aatatgtata	ttaaaaggaa	aatcacactt	taaaaatgtg	aggaaagctt	tgaaaacagt	600
cttaatgcat	gagtcacatc	acatattttc	aagttttgga	aacagaaaga	agtttagaat	660
tttcaaagta	atctgaaaac	tttctaagcc	attttaaaat	aagatttttt	tccccatctt	720
tccaatgttt	cctatttgat	agtgtaatat	agaaatgggc	agtttctagt	gtcgaacttaa	780
ctgtgctaata	tcataagtca	ttatacattt	atgactttaag	agttcaaata	agtggaaatt	840
gggttataat	gaaaatgaca	agggggcccc	ttcagcagcc	actcatctga	actagtaatc	900
ccaacacaat	g					911
<210> 92						
<211> 1710						
<212> DNA						
<213> Homo sapiens						
<400> 92						
tttttttttt	tttttaactt	ttagcagtg	ttatttttgt	taaaagaaac	caattgaatt	60
gaaggtcaag	acaccttctg	attgcacaga	ttaaacaaga	aagtattact	tatttcaact	120
ttacaaagca	tcttattgat	ttaaaaagat	ccatactatt	gataaagttc	accatgaaca	180
tatatgtaat	aaggagacta	aaatattcat	tttaccatct	tacaacatgt	atttcatatt	240
tctaataaac	cacaaatcat	ataggaaaat	atttaggtcc	atgaaaaagt	ttcaaaacat	300
taaaaaatta	aagttttgaa	acaaatcaca	tgtgaaagct	cattaaataa	taacattgac	360
aaataaatag	ttaatcagct	ttacttatta	gctgtgcca	tgcatttctg	gcattccatt	420
ccaagcgagg	gtcagcatgc	aggggtataat	ttcatactat	gcgaccgtaa	agagctacag	480
ggcttatttt	tgaagtgaag	tgtcacaggg	tctttcatte	tctttcaaag	gaagatcact	540
catggctgct	aaactgttcc	catgaagagt	acaaaaaag	cacctttctg	aaatgttact	600
gtgaagattc	atgacaacat	atttttttta	acctgttttg	aaggagtttt	gtttaggaga	660
ggggatgggc	cagtagatgg	aggggtatctg	agaagccctt	ttctgtttta	aaatataatg	720
attcactgat	gtttatagta	tcaacagctc	tttaagaaca	atgaggaatt	aaaactacag	780
gatactgga	atttaaatgc	aaattgcatt	catggatata	cctacatctt	gaaaaacttg	840
aaaaggaaaa	actattccca	aagaaggtcc	tgatacttaa	gacagcttgc	tgggtttgat	900
caaagcagaa	agcatatact	ttcaagtga	aaaacagcag	tggcaggctt	gagctttcca	960
agcaatcaaa	tctgtaaagc	agatggttac	tagtaagtct	agttatggga	gtctgagttc	1020
taactcatgc	tgtgcttgct	ggatttgctg	gctcttttcc	gctctctgtg	atgctggact	1080
ggcttgctag	gtgacatgct	ctcaaagttg	tgactggact	cgttgtgctg	ccgggtgtac	1140
ctcttgact	tgcaggcagt	gactactgtg	attttgtagg	tgcgtgtgct	gccatcttgg	1200
cactgcagct	ggattctctg	ggtacgggtt	ttgtcattga	cacaccgcca	ctcctgggag	1260
ctcctcctgc	tccagtactt	tgttccatag	cctcctccaa	tccagttagg	gagcactggc	1320
aggggcaagc	actcgccagc	acacaccagc	tccttcagag	ggctgatgct	ggtgcactgg	1380
ccatcagaga	tgtatttgg	ggaacgcagt	tcccggcaat	ccacttgaa	ccgagtgttc	1440
cgatccagtc	cagtgttact	gaaatgcctg	cctccatttc	tggcttgatt	caacgtgctg	1500
ttgtgctg	ggtgtgctg	aacaggttta	accacatgtg	aataaaggat	ttctgtggca	1560
tcatttttaa	aagccaaaca	gcttttcatt	aggatgcatg	caaggggaag	gagatagaaa	1620
tgaatggcag	gaggaagcat	ggtgagtaga	ggatttgctt	gactgaagag	ctggttaatt	1680
cttttgctc	tgcccaacac	aatggaattc				1710

&lt;210&gt; 93

&lt;211&gt; 251

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 93

cccaccctac ccaaataatta gacaccaaca cagaaaagct agcaatggat tcccttctac 60  
tttgttaaat aaataagtta aatatttaaa tgctgtgtgc tctgtgatgg caacagaagg 120  
accaacaggc cacatcctga taaaaggtaa gaggggggtg gatcagcaaa aagacagtgc 180  
tgtgggctga ggggacctgg ttcttgtgtg ttgccctca agactcttcc cctacaaata 240  
actttcatat g 251

&lt;210&gt; 94

&lt;211&gt; 738

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 94

cccttttttt ttttttttcc acttctcagt ttatttctgg gactaaattt gggctcagagc 60  
tgcagagaag ggatgggccc tgagcttgag gatgaaagtg ccccaggag attgagagcg 120  
aaccctcgcc ctggacagtt ttggaaattg ttcccagggt tcaactagag agacacggtc 180  
agcccaatgt gggggaagca gacctgagt ccaggagaca tggggtcagg ggctggagag 240  
atgaacattc tcaacatctc tgggaaggaa tgagggtctg aaaggagtgt cagggtctgc 300  
cctgcagcag gtggggatgc cgggtgtgctg agtcctggga tgactcagga gttggcctgg 360  
atggtttcct ggatccactt ggtgaacttg cagaggttcg ttagacacc cggctctgtg 420  
ggccggggcag aagggtaatc tcccaggac acgagtcctt gcaggagacc attgcagacc 480  
acaggccccc cagaatcacc ctggcaggag tctctacctg ctttgtcacc ggcgcagaac 540  
atggtgtcat ctatctgtct cgggtaagca tctcgcacc ttttctgact tagcacgtg 600  
atattcaagc actggaggac cttaggaag tgacttggg ggctcttggg tgccccccag 660  
ccagacacca agcactttgt cccagcagag ggacaatgag aggagacgtt gatgggtctg 720  
acatctttag tgggacga 738